## FLI1 Monoclonal Mouse Antibody (FLI1/1312)

## **Product Description**

This antibody recognizes a protein of 51 kDa, which is identified as FLI1. This protein, a member of the ETS family of DNA binding transcription factors, is involved in cellular proliferation and tumorigenesis. Ets-1 is the prototype member of a family of genes identified on the basis of homology to the v-Ets oncogene isolated from the E26 erythroblastosis virus. Members of the Ets gene family share a highly conserved carboxy-terminal domain containing a sequence related to the SV40 large T antigen nuclear localization signal sequence. Approximately 90% of Ewing s Sarcoma (EWS)/Primitive Neuroectodermal Tumors (PNET) have a specific translocation, t(11;22)(q24;q12), which results in fusion of EWS to Fli-1, and production of an EWS-Fli-1 fusion protein. Among normal tissues only endothelial cells and small lymphocytes express Fli-1. This protein is expressed in majority of vascular tumors including angiosarcomas, hemangioendotheliomas, hemangiomas, and Kaposi's Sarcomas. High sensitivity and specificity of Fli-1 equals to or exceeds that of the established vascular markers like CD31, CD34, and Factor VIII. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the <u>CF® Dye Brochure</u> for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 1312, Anti-FLI1 (FLI1/1312)

## **Product attributes**

Call us: 800-304-5357 Email: btinfo@biotium.com

Product attributes	•			
Antibody number	#1312			
Antibody reactivity	FLI1			
(target) Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	FLI1/1312			
Isotype	IgG			
Molecular weight	51 kDa			
Synonyms	ERGB transcription factor; Ewing Sarcoma breakpoint region 2 (EWSR2); FL11; FL11 EWS fusion gene; Friend leukemia integration 1 (FL11) transcription factor; Friend leukemia virus integration 1; Proto-nocogene Fli-1 SIC1; Transcription factor ERGB; Viral integration region FL11			
Human gene symbol	FLI1			
Entrez gene ID	2313			
SwissProt	Q01543			
Unigene	504281			
Immunogen	Recombinant fragment (aa 107-261) of human FLI1 protein (exact sequence is proprietary)			
Verified antibody	WB (verified)			
Antibody target cellular	WB (verified) Nucleus			
annlications	, ,			
Antibody target cellular	Nucleus Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL,			
Antibody target cellular localization Species reactivity  Antibody application	Nucleus Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optime dilution for a specific application should			
Antibody target cellular localization Species reactivity Antibody application notes	Nucleus Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optima dilution for a specific application should be determined by user THP1 and Raji cells. Tonsil, Ewing s			
Antibody target cellular localization Species reactivity Antibody application notes  Positive control	Nucleus Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optima dilution for a specific application should be determined by user THP1 and Raji cells. Tonsil, Ewing s sarcoma (EWS) and Angiosarcoma			
Antibody target cellular localization Species reactivity Antibody application notes  Positive control Shipping condition	Nucleus Human  Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optima dilution for a specific application should be determined by user  THP1 and Raji cells. Tonsil, Ewing s sarcoma (EWS) and Angiosarcoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store			
Antibody target cellular localization Species reactivity Antibody application notes  Positive control Shipping condition Storage Conditions	Nucleus Human  Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optime dilution for a specific application should be determined by user  THP1 and Raji cells. Tonsil, Ewing s sarcoma (EWS) and Angiosarcoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as			
Antibody target cellular localization Species reactivity Antibody application notes  Positive control Shipping condition Storage Conditions  Shelf life	Nucleus Human  Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optima dilution for a specific application should be determined by user  THP1 and Raji cells. Tonsil, Ewing s sarcoma (EWS) and Angiosarcoma Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2			
Antibody target cellular localization Species reactivity Antibody application notes  Positive control Shipping condition Storage Conditions  Shelf life  Regulatory status Antibody/conjugate	Nucleus Human  Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optima dilution for a specific application should be determined by user  THP1 and Raji cells. Tonsil, Ewing s sarcoma (EWS) and Angiosarcoma Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide Purified, BSA-free: 1 mg/mL in PBS			
Antibody target cellular localization Species reactivity  Antibody application notes  Positive control  Shipping condition  Storage Conditions  Shelf life  Regulatory status  Antibody/conjugate formulation	Nucleus Human  Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optima dilution for a specific application should be determined by user  THP1 and Raji cells. Tonsil, Ewing s sarcoma (EWS) and Angiosarcoma Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purffied: 0.2 mg/mL in PBS/0.05% BSA, Purffied: 0.5 without azide Cancer, Immunology, Transcription			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	<b>Detection channel</b>	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.