VLDL Receptor Monoclonal Mouse Antibody (VLDLR/1337)

Product Description

VLDLR (very low density lipoprotein receptor) is a member of the LDL receptor gene family, which includes LDL receptor, LRP, megalin, VLDLR and ApoER2. The LDL receptor family is characterized by a cluster of cysteine-rich class A repeats, epidermal growth factor (EGF)-like repeats, YWTD repeats and an O-linked sugar domain. VLDLR associates with RAP (receptor associated protein) during receptor folding, and RAP facilitates the secretion of the extracellular region of VLDLR. VLDLR is thought to mediate the interaction of extracellular Reelin and cytosolic mDab1 (mammalian disabled protein), which activates a tyrosine kinase. This pathway regulates the migration of neurons along the radial glial fiber network during brain development.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1337, Anti-VLDL Receptor (VLDLR/1337)

Product attributes

Call us: 800-304-5357

Product attributes				
Antibody number	#1337			
Antibody reactivity (target)	VLDL Receptor			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	VLDLR/1337			
Isotype	lgG1			
Molecular weight	143-161 kDa			
Synonyms	Very low-density lipoprotein receptor; VLDL receptor; VLDL-R; VLDLR			
Human gene symbol	VLDLR			
Entrez gene ID	7436 (Human)			
SwissProt	P98155 (Human)			
Unigene	370422 (Human)			
Immunogen	Recombinant human VLDLR fragment from c-terminal (exact sequence is proprietary)			
Verified antibody applications	IHC (FFPE) (verified)			
Antibody target cellular localization	Plasma membrane			
Species reactivity	Human, Rat			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology (formalin) 1-2 ug/mL, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user			
Antibody application notes Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology (formalin) 1-2 ug/mL, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western			
	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology (formalin) 1-2 ug/mL, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal diultion for a specific application should be determined by user HeLa or U-251 cells. Pancreas, Cerebellum, Heart or Skeletal			
Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology (formalin) 1-2 ug/mL, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HeLa or U-251 cells. Pancreas, Cerebellum, Heart or Skeletal Muscle.			
Positive control Shipping condition	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Immunofluorescence: 0.5-1 ug/ml., Immunofluology (formalin) 1-2 ug/ml., Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/mllillon cells/0.1 ml., Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should be determined by user HeLa or U-251 cells. Pancreas, Cerebellum, Heart or Skeletal Muscle. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Positive control Shipping condition Storage Conditions	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology (formalin) 1-2 ug/mL, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should be determined by user HeLa or U-251 cells. Pancreas, Cerebellum, Heart or Skeletal Muscle. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when			
Positive control Shipping condition Storage Conditions Shelf life	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology (formalin) 1-2 ug/mL, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HeLa or U-251 cells. Pancreas, Cerebellum, Heart or Skeletal Muscle. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			
Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology (formalin) 1-2 ug/mL, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM Tris, 1 mM EDTA pH 9.0 for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HeLa or U-251 cells. Pancreas, Cerebellum, Heart or Skeletal Muscle. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 1 mg/mL in PBS/0.05% BSA, Purified: 1 mg/mL in PBS/0.05% BSA, Purified: 1 mg/mL in PBS/0.05% ESA, Purified: 1 mg/mL			

Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on July 12, 2025 at 02:47:26 AM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/mouse-monoclonal-anti-vldl-receptor-vldlr1337/