## NSE-Gamma Monoclonal Mouse Antibody (ENO2/1375)

## **Product Description**

This antibody recognizes a protein of about 50 kDa, which is identified as gamma-enolase. Three isoenzymes of enolases are identified, alpha, beta and gamma. Alpha-isoform is expressed in most tissues, whereas beta-form is expressed predominantly in muscle tissue whereas gamma-enolase is found only in nervous tissue. These isoforms exist as both homodimers and heterodimers, and they play a role in converting phosphoglyceric acid to phosphenolpyruvic acid in the glycolytic pathway. NSE-gamma is a useful marker to identify peripheral nerves and tumors of neuro-endocrine origins, such as pheochromocytomas. It it be usually employed in combination with other markers such as Synaptophysin, Chromogranin A, and Neurofilament.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

**Stock status:** Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1375, Anti-NSE-Gamma (ENO2/1375)

## Oradust attributes

Call us: 800-304-5357

Product attributes				
Antibody number	#1375			
Antibody reactivity (target)	NSE-Gamma			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	ENO2/1375			
Isotype	lgG2b			
Molecular weight	~50 kDa			
Synonyms	2-phospho-D-glycerate hydrolyase; ENO2; ENOG; Enolase 2 gamma neuronal; Enolase2; Gamma-enolase; Neural enolase Neuron specific gamma enolase; Neuron-specific enolase; NS			
Human gene symbol	ENO2			
Entrez gene ID	2026			
SwissProt	P09104			
Unigene	511915			
Immunogen	A synthetic peptide corresponding to aa416-433 of human NSE gamma (exact sequence is proprietary)			
Antibody target cellular localization	Cytoplasmic			
Verified antibody applications	IHC (FFPE) (verified), WB (verified)			
Species reactivity	Human, Mouse, Rat			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology (formalin) 0.1-0.2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/mlilion cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user			
	blotting 0.5-1 ug/mL, Optimal dilution for a specific application			
Positive control	blotting 0.5-1 ug/mL, Optimal dilution for a specific application			
Positive control Shipping condition	blotting 0.5-1 ug/mL, Óptimal dilution for a specific application should be determined by user  HepG2, SH-SY-5Y, HeLa or Y79 cells. Pancreas, Cerebellum or			
	blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user  HepG2, SH-SY-5Y, HeLa or Y79 cells. Pancreas, Cerebellum or Pheochromocytoma.			
Shipping condition	blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HepG2, SH-SY-5Y, HeLa or Y79 cells. Pancreas, Cerebellum or Pheochromocytoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Shipping condition Storage Conditions	blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user  HepG2, SH-SY-5Y, HeLa or Y79 cells. Pancreas, Cerebellum or Pheochromocytoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when			
Shipping condition Storage Conditions Shelf life	blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user  HepG2, SH-SY-5Y, HeLa or Y79 cells. Pancreas, Cerebellum or Pheochromocytoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user  HepG2, SH-SY-5Y, HeLa or Y79 cells. Pancreas, Cerebellum or Pheochromocytoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user  HepG2, SH-SY-5Y, HeLa or Y79 cells. Pancreas, Cerebellum or Pheochromocytoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purfiled. BSA-free: 1 mg/mL in PBS without azide			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Antibody research areas	blotting 0.5-1 uý/mL, Optimal dilution for a specific application should be determined by user  HepG2, SH-SY-5Y, HeLa or Y79 cells. Pancreas, Cerebellum or Pheochromocytoma.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide  Cancer, Metabolism, Neuroscience  Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in			

Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on September 10, 2025 at 06:06:32 PM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/mouse-monoclonal-anti-nse-gamma-eno21375/">https://biotium.com/product/mouse-monoclonal-anti-nse-gamma-eno21375/</a>