Alpha-1 Antichymotrypsin Monoclonal Mouse Antibody (AACT/1451 + AACT/1452)

Product Description

Alpha-1 Antichymotrypsin (AACT) is a plasma protease inhibitor synthesized in the liver as a single glycopeptide chain. In human, the normal serum level of AACT is about one-tenth that of alpha-1-antitrypsin (AAT), with which it shares nucleic acid and protein sequence homology. Both are major acute phase reactants; their concentrations in plasma increase in response to trauma, surgery and infection. Elevated levels of AACT are widely, but not universally, reported in the cerebrospinal fluid and plasma of AD patients. Prostate-specific antigen (PSA) and its SDS-stable complex with AACT are in widespread use as markers for the diagnosis of prostate cancer. AACT deficiency may also be a possible cause of chronic liver disease. AACT antibody reacts with histiocytes and histiocytic neoplasms. It is widely used to identify histiocytes and tumors derived from them. Acinar tumors of the pancreas and salivary gland may also exhibit AACT positivity.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1453, Anti-Alpha-1 Antichymotrypsin (AACT/1451 AACT/1452)

Product attributes

Call us: 800-304-5357

Product attributes			
Antibody number	#1453		
research-areas	Cancer		
Antibody reactivity (target)	Alpha-1 Antichymotrypsin		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	AACT/1451 + AACT/1452		
Isotype	lgG1		
Molecular weight	65-76 kDa		
Synonyms	Alpha-1-antichymotrypsin; Serpin A3; SERPINA3; AACT; ACT; Cell growth-inhibiting gene 24/25 protein; Growth inhibiting protein 24; Growth inhibiting protein 25; Serine (or cysteine) proteinase inhibitor clade A member 3		
Human gene symbol	SERPINA3		
Entrez gene ID	12		
SwissProt	P01011		
Unigene	534293; 710488		
Immunogen	Recombinant human Antichymotrypsin (AACT) protein fragment (aa49-187) (exact sequence is proprietary)		
Verified antibody applications	IHC (FFPE) (verified)		
Antibody target cellular localization	Secreted (extracellular)		
	Human		
Species reactivity	Human		
Species reactivity Antibody application notes	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/ml. Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mlilion cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mllilion cells/0.1 mL, Optimal dilution for a specific application should be		
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mllilion cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
Antibody application notes Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mLl Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Optimal dilution for a specific application should be determined by user HeLa Cells. Tonsil, Pancreas or Histiocytoma.		
Antibody application notes Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user HeLa Cells. Tonsil, Pancreas or Histiocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
Antibody application notes Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mllilion cells/0.1 mL, Optimal dilution for a specific application should be determined by user HeLa Cells. Tonsil, Pancreas or Histiocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when		
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mllilion cells/0.1 mL, Optimal dilution for a specific application should be determined by user HeLa Cells. Tonsil, Pancreas or Histiocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended		
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Optimal dilution for a specific application should be determined by user HeLa Cells. Tonsil, Pancreas or Histiocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purffied .0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% azide, Purffied .0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.0		
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mllilion cells/0.1 mL, Optimal dilution for a specific application should be determined by user HeLa Cells. Tonsil, Pancreas or Histiocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/1.05% azide, Purffied: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purffied: 0.2 mg/mL in PBS/		
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Antibody research areas	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mllillion cells/0.1 mL, Optimal dilution for a specific application should be determined by user HeLa Cells. Tonsil, Pancreas or Histiocytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% BSA/ree: 1 mg/mL in PBS without azide Cancer Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in		
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Antibody research areas Product origin	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Immunofluorescence 0.5-1 ug/mL Western blotting 0.5-1 ug/mL, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Optimal dilution for a specific application should be determined by user HeLa Cells. Tonsil, Pancreas or Histicoytoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Cancer Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot.		

Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on September 10, 2025 at 10:45:12 AM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/mouse-monoclonal-anti-alpha-1-antichymotrypsin-aact1451-aact1452/