## CD80 Monoclonal Mouse Antibody (C80/1608)

## **Product Description**

T cell proliferation and lymphokine production are triggered by occupation of the TCR by antigen, followed by a costimulatory signal that is delivered by a ligand expressed on antigen presenting cells. The B7-related cell surface proteins CD80 (B7-1) and CD86 (B7-2) are expressed on antigen presenting cells and bind the homologous T cell receptors CTLA-4 (cytotoxic T lymphocyte-associated protein-4) and CD28 and trigger costimulatory signals for optimal T cell activation. CTLA-4 shares 31% overall amino acid identity with CD28 and it has been proposed that CD28 and CTLA-4 are functionally redundant. SLAM is a novel receptor on T cells that, when engaged, potentiates T cell expansion in a CD28-independent manner. B7, also designated BB1, is another ligand or counter receptor for CD28 and CTLA-4 that is expressed on the antigen-presenting cell. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and See the <u>CF® Dye Brochure</u> for more information. Note: Conjugates of blue inderescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. **Stock status**: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 1608, Anti-CD80 (C80/1608)

Call us: 800-304-5357 Email: btinfo@biotium.com

Product attributes				
Antibody number	#1608			
Antibody reactivity	CD80			
(target) Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	C80/1608			
Isotype	IgG1, kappa			
Molecular weight	60 kDa			
Synonyms	Activation B7-1 antigen; B lymphocyte activation antigen B7; B7-1; BB1; CD28 antigen ligand 1; CD28LG; CD28LG1; CD80; Costimulatory factor CD80; CTLA-4 counter-receptor B7.1; LAB7; T-lymphocyte activation antigen CD80			
Human gene symbol	CD40			
Entrez gene ID	941			
SwissProt	P33681			
Unigene	838			
Immunogen	Recombinant human full-length CD80 protein			
Antibody target cellular	Plasma membrane			
Species reactivity	Human			
Antibody application notes	For coating for ELISA, order Ab without BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user. Recommended starting concentrations for titration are 1-2 ugml. for most applications, or 1 ugmillion cells/100 ut for flow cytometry			
	indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL			
Positive control	indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL			
Positive control Shipping condition	indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user. Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry  Raji or Ramos Cells. Tonsil, Spleen or			
	indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry  Raji or Ramos Cells. Tonsil, Spleen or Thymus  Room temperature  Store at 2 to 8 °C, Protect fluorescent			
Shipping condition	indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user. Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry  Raji or Ramos Cells. Tonsil, Spleen or Thymus  Room temperature			
Shipping condition	indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user. Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry  Raji or Ramos Cells. Tonsil, Spleen or Thymus  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store			
Shipping condition Storage Conditions	indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml. for most applications, or 1 ug/million cells/100 uL for flow cytometry  Raji or Ramos Cells. Tonsil, Spleen or Thymus  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as			
Shipping condition Storage Conditions Shelf life	indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry  Raji or Ramos Cells. Tonsil, Spleen or Thymus  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry Raji or Ramos Cells. Tonsil, Spleen or Thymus Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Free: 1 mg/mL in PBS/0.05% azide, Purified; 0.2 mg/mL in PBS/0.05% BSA-free: 1 mg/mL in PBS/0.05% azide, Purified; 0.84-free: 1 mg/mL in PBS/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.