Connexin 32 Monoclonal Rat Antibody (R5.21C)

Product Description

This antibody recognizes a protein of 27-32 kDa, identified as Connexin 32. The connexin family of proteins forms hexameric complexes called connexons that facilitate movement of low molecular weight proteins between cells via gap junctions. Connexin proteins share a common topology of four transmembrane alphalpha-helicaldomains, two extracellular loops, a cytoplasmic loop and cytoplasmic N- and C-termini. Many of the key functional differences arise from specific amino-acid substitutions in the most highly conserved domains, the transmembrane and extracellular regions. Each of the approximately 20-connexin isoforms produces channels with distinct permeability and electrical and chemical sensitivities; therefore, one connexin usually cannot fully substitute for another. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the <u>CF® Dye Brochure</u> for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 1641, Anti-Connexin-32 (R5.21C)

References

Note: References for this clone sold by other suppliers may be listed for expected applications.

- 1. J Cell Biol (1986) 103 (3): 755-766. (WB)
- Endocrinology (1993) 133: 2371-2378. (IF)
 J Cell Sci (1994) 107: 955-967. (IP)
- 4. Int J Cancer (1997) 73: 479-485. (IHC, frozen)

BSA-free

Call us: 800-304-5357

Product attributes				
Antibody number	#1641			
Antibody reactivity	Connexin-32			
(target) Antibody type	Primary			
Host species	Rat			
Clonality	Monoclonal			
Clone	R5.21C			
Isotype	lgG2a			
Molecular weight	27-32 kDa			
Synonyms	Charcot Marie Tooth neuropathy X linked; CMTX; CMTX1; Connexin-32; Cx32; GAP junction 28kDa liver protein; Gap junction beta-1 protein; Gap junction protein beta 1 32kD; GJB1			
Human gene symbol	Gjb1			
Entrez gene ID	14618 (Mouse)			
SwissProt	P28230 (Mouse)			
Unigene	21198 (Mouse)			
Immunogen	Mouse liver DOC-JR-plasma membranes			
Antibody target cellular Species reactivity	Plasma membrane			
	Mouse, Rat			
Expected antibody applications	IHC (frozen) (published for clone), IF (published for clone), IP (published for clone), WB (published for clone)			
	cione), wb (published for cione)			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Immunofistology (frozen) 0.5-1 ug/mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user			
	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/mllion cells, Immunohistology (frozen) 0.5-1 ug/ml. Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should			
notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunofluorescence: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, immunohistology (frozen) 0.5-1 ug/mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user MCF-7 Cells. Pancreas, Liver, Kidney,			
Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunofluorescence: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, immunohistology (frozen) 0.5-1 ug/mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user MCF-7 Cells. Pancreas, Liver, Kidney, Stomach or Tonsil			
Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Immunohistology (frozen) 0.5-1 ug/mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user MCF-7 Cells. Pancreas, Liver, Kidney, Stomach or Tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store			
Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Inmunofluorescence: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Immunofluorescence: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Immunofluorescence: 1-2 ug/mL, Optimal dilution for a specific application should be determined by user MCF-7 Cells. Pancreas, Liver, Kidney, Stomach or Tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as			
Positive control Shipping condition Storage Conditions Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Interest that the conjugates than for indirect detection with secondary antibody. Immunofluorescence: 1-2 ug/mL, Flow cytometry: 0.5-1 ug/million cells, Immunohistology (frozen) 0.5-1 ug/mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user MCF-7 Cells. Pancreas, Liver, Kidney, Stomach or Tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			

Email: btinfo@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.