## von Willebrand Factor / vWF Monoclonal Mouse Antibody (F8/86)

## **Product Description**

von Willebrand Factor (vWF) is a multimeric glycoprotein that is found in endothelial cells, plasma and platelets. It acts as a carrier protein for Factor VIII and promotes platelet adhesion and aggregation. vWF undergoes a variety of posttranslational modifications that influence the affinity and availability for Factor VIII, including cleavage of the propeptide and formation of N-terminal disulfide bonds. This antibody helps to establish the endothelial nature of some lesions of disputed histogenesis, e.g. Kaposi's sarcoma and cardiac myxoma. It is widely used for differentiating vascular lesions from those of other tissue differentiation within a panel of other vascular markers although not all tumors of endothelial differentiation contain this antigen.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1716, Anti-von Willebrand Factor (F8/86)

## Oradust attributes

| Product attributes                    |   |  |  |
|---------------------------------------|---|--|--|
| Antibody number                       | #1716   |  |  |
| Antibody reactivity (target)          | von Willebrand Factor, vWF  |  |  |
| Antibody type                         | Primary   |  |  |
| Host species                          | Mouse   |  |  |
| Clonality                             | Monoclonal  |  |  |
| Clone                                 | F8/86   |  |  |
| Isotype                               | lgG1  |  |  |
| Molecular weight                      | 250 kDa   |  |  |
| Synonyms                              | Coagulation Factor VIII; Factor VIII Related Antigen; F8VWF; von Willebr; Antigen 2; von Willebr; Disease (vWD)   |  |  |
| Human gene symbol                     | VWF   |  |  |
| Entrez gene ID                        | 7450  |  |  |
| SwissProt                             | P04275  |  |  |
| Unigene                               | 440848  |  |  |
| Immunogen                             | Factor isolated from human plasma, von Willebr  |  |  |
| Antibody target cellular localization | Secreted (extracellular), Vesicular   |  |  |
| Verified antibody applications        | IHC (FFPE) (verified)   |  |  |
| Species reactivity                    | Human   |  |  |
| Antibody application notes            | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology (formalin) 1-2 ug/mL, Staining of formalin-fixe tissues requires boiling tissue sections in 10 mM citrate buffer pH 6.0, for 10-20 min followed by cooling at RT for 20 min, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user |  |  |
|                                       | specific application should be determined by user   |  |  |
| Positive control                      | specific application should be determined by user  HUVEC cells or Tonsil  |  |  |
| Positive control Shipping condition   | · · · ·   |  |  |
|                                       | HUVEC cells or Tonsil   |  |  |
| Shipping condition                    | HUVEC cells or Tonsil  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,   |  |  |

Call us: 800-304-5357 Email: techsupport@biotium.com

|                                | 110 120 00110 01 1011011  |  |  |
|--------------------------------|---|--|--|
| Shipping condition             | Room temperature  |  |  |
| Storage Conditions             | Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C   |  |  |
| Shelf life                     | Guaranteed for at least 24 months from date of receipt when stored as recommended   |  |  |
| Regulatory status              | For research use only (RUO)   |  |  |
| Antibody/conjugate formulation | Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide |  |  |
| Antibody research areas        | Hematology  |  |  |
| Product origin                 | Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot.                   |  |  |
| Cell/tissue expression         | Endothelial cells   |  |  |

| Antibody # prefix | Conjugation           | Ex/Em (nm) | Laser line | Detection channel           | Dye Features     |
|-------------------|-----------------------|------------|------------|-----------------------------|------------------|
| BNC04             | CF®405S               | 404/431    | 405        | DAPI (microscopy),<br>AF405 | CF®405S Features |
| BNC88             | CF®488A               | 490/515    | 488        | GFP, FITC                   | CF®488A Features |
| BNC68             | CF®568                | 562/583    | 532, 561   | RFP, TRITC                  | CF®568 Features  |
| BNC94             | CF®594                | 593/614    | 561        | Texas Red®                  | CF®594 Features  |
| BNC40             | CF®640R               | 642/662    | 633-640    | Cy®5                        | CF®640R Features |
| BNC47             | CF®647                | 650/665    | 633-640    | Cy®5                        | CF®647 Features  |
| BNC74             | CF®740                | 742/767    | 633-685    | 775/50                      | CF®740 Features  |
| BNCB              | Biotin                | N/A        | N/A        | N/A                         |                  |
| BNUB              | Purified              | N/A        | N/A        | N/A                         |                  |
| BNUM              | Purified,<br>BSA-free | N/A        | N/A        | N/A                         |                  |

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on July 12, 2025 at 07:23:05 AM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/mouse-monoclonal-anti-von-willebr-factor-factor-viii-related-ag-f886/">https://biotium.com/product/mouse-monoclonal-anti-von-willebr-factor-factor-viii-related-ag-f886/</a>