
biotium Glowing products for science

Call us : 800-304-5357

# CD269 / TNFRSF17 / BCMA Monoclonal Mouse Antibody (BCMA/2366)



## Product Description

The B cell maturation protein (BCMA) is a type I integral membrane protein that belongs to the tumor necrosis factor receptor (TNF-R) superfamily. It is expressed as a 184 amino acid peptide that is expressed only in mature B-lymphocytes and is located on the cis part of the Golgi apparatus. BCMA shares significant homology with TACI (transmembrane activator) within the cysteine-rich domain. TACI has been shown to bind CAML, which induces activation of NFAT (nuclear factor of activated T cells). Both BCMA and TACI have been shown to bind APRIL and TALL-1, which stimulate B cell proliferation in conjunction with other B-cell activators. When overexpressed, TALL-1 stimulates the development of systemic lupus erythaematosus (SLE).

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

**Stock status:** Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

**Catalog number key for antibody number 2366, Anti-CD269 / TNFRSF17 / BCMA (BCMA/2366)**

## Product attributes

| | |
|---|---|
| **Antibody number** | #2366 |
| **Antibody reactivity (target)** | BCMA, CD269, TNFRSF17 |
| **Antibody type** | Primary |
| **Host species** | Mouse |
| **Clonality** | Monoclonal |
| **Clone** | BCMA/2366 |
| **Isotype** | IgG2c, kappa |
| **Molecular weight of antigen** | 20 kDa |
| **Synonyms** | B-cell maturation protein; BCMA; CD269; TNR17; Tumor necrosis factor receptor superfamily member 17 (TNFRSF17) |
| **Human gene symbol** | CD269 |
| **Entrez gene ID** | 608 |
| **SwissProt** | Q02223 |
| **Unigene** | 2556 |
| **Immunogen** | Recombinant human CD269 protein fragment (around aa 78-184) (exact sequence is proprietary) |
| **Verified antibody applications** | IHC (FFPE) (verified) |
| **Antibody target cellular localization** | Plasma membrane |
| **Species reactivity** | Human |
| **Antibody application notes** | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, ELISA: For coating order antibody without BSA, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user |
| **Positive control** | NAMALWA Cells. Tonsil. Gastric Carcinoma |
| **Shipping condition** | Room temperature |
| **Storage Conditions** | Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C |
| **Shelf life** | Guaranteed for at least 24 months from date of receipt when stored as recommended |
| **Regulatory status** | For research use only (RUO) |
| **Antibody/conjugate formulation** | Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide |
| **Validated in protein array** | Monospecific |
| **Antibody research areas** | Autoimmunity, Immunology |
| **Product origin** | Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot. |
| **Cell/tissue expression** | B-cells |

| Antibody # prefix | Conjugation | Ex/Em (nm) | Laser line | Detection channel | Dye Features |
|---|---|---|---|---|---|
| BNC04 | CF®405S | 404/431 | 405 | DAPI (microscopy), AF405 | CF®405S Features |
| BNC88 | CF®488A | 490/515 | 488 | GFP, FITC | CF®488A Features |
| BNC68 | CF®568 | 562/583 | 532, 561 | RFP, TRITC | CF®568 Features |
| BNC94 | CF®594 | 593/614 | 561 | Texas Red® | CF®594 Features |
| BNC40 | CF®640R | 642/662 | 633-640 | Cy®5 | CF®640R Features |
| BNC47 | CF®647 | 650/665 | 633-640 | Cy®5 | CF®647 Features |
| BNC74 | CF®740 | 742/767 | 633-685 | 775/50 | CF®740 Features |
| BNCB | Biotin | N/A | N/A | N/A | |
| BNUB | Purified | N/A | N/A | N/A | |
| BNUM | Purified, BSA-free | N/A | N/A | N/A | |

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.