
CD79b Monoclonal Mouse Antibody (IGB/1844)



Call us: 800-304-5357

Product Description

CD79 (also designated Ig α /Ig β) is a heterodimer composed of α chains, designated CD79A or MB-1, and β chains, designated CD79B or B29. The B cell antigen receptor complex (BCR) is formed by the association of CD79 with a membrane immunoglobulin, such as IgM or IgD. The membrane immunoglobulins IgM and IgD achieve surface expression and antigen presentation function in response to CD79 association. The cytoplasmic tails of both CD79A and CD79B contain an ITAM (immuno-receptor tyrosine-based activation) motif, which acts to initiate the BCR signaling reactions by binding to and activating tyrosine kinases. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 1844, Anti-CD79b (IGB/1844)



Positive control Shipping condition

Storage Conditions

Regulatory status

formulation

Antibody/conjugate

Validated in protein

Tumor expression

array Antibody research areas Immunology Cell/tissue expression NK cells, T-cells

Shelf life

Email: btinfo@biotium.com

Daudi or Ramos cells. Germinal center B-cells in a lymph node or tonsil.

Store at 2 to 8 °C, Protect fluorescent

conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C

For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1%

Guaranteed for at least 24 months from date of receipt when stored as recommended

Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide Purified, BSA-free: 1 mg/mL in PBS without azide

Room temperature

Monospecific

Leukemia/lymphoma

| Antibody # prefix BNC04 | Conjugation CF®405S | Ex/Em (nm) 404/431 | Laser line 405 | Detection channel DAPI (microscopy), AF405 | Dye Features CF®405S Features |
|----------------------------|------------------------|-----------------------|-------------------|--|---|
| BNC88 | CF®488A | 490/515 | 488 | GFP, FITC | CF®488A Features |
| BNC68 | CF®568 | 562/583 | 532, 561 | RFP, TRITC | CF®568 Features |
| BNC94 | CF®594 | 593/614 | 561 | Texas Red® | CF®594 Features |
| BNC40 | CF®640R | 642/662 | 633-640 | Cy®5 | CF®640R Features |
| BNC47 | CF®647 | 650/665 | 633-640 | Cy®5 | CF®647 Features |
| BNCB | Biotin | N/A | N/A | N/A | |
| BNUB | Purified | N/A | N/A | N/A | |
| BNUM | Purified, | N/A | N/A | N/A | |

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience

BSA-free