## Interleukin 10 Recombinant Monoclonal Rabbit Antibody (IL10/2651R)



## **Product Description**

Interleukin 10 (IL10) is a cytokine produced primarily by monocytes and to a lesser extent by lymphocytes. This cytokine has pleiotropic effects in immuno-regulation and inflammation. It down-regulates the expression of Th1 cytokines, MHC class II antigens, and costimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. This cytokine can block NF-kappa B activity, and is involved in the regulation of the JAK-STAT signaling pathway. Knockout studies in mice suggested the function of this cytokine as an essential immuno-regulator in the intestinal tract. Mutations in this gene are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 2651, Anti-Interleukin 10 (IL10/2651R)

Call us: 800-304-5357

Product attributes				
Antibody number	#2651			
Antibody reactivity (target)	Interleukin 10			
Antibody type	Primary			
Host species	Rabbit			
Clonality	Recombinant Monoclonal			
Clone	IL10/2651R			
Isotype	IgG			
Molecular weight	20 kDa (Monomer); 37 kDa (Dimer)			
Synonyms	Cytokine Synthesis Inhibitory Factor (CSIF); GVHDS; IL-10; IL10; IL10A; Interleukin-10; T-cell Growth Inhibitory Factor (TGIF)			
Human gene symbol	IL10			
Entrez gene ID	3586			
SwissProt	P22301			
Unigene	193717			
Immunogen	Recombinant full-length human Interleukin 10 (IL10) protein			
Verified antibody applications	IHC (FFPE) (verified)			
Antibody target cellular localization	Jular Secreted (extracellular)			
Species reactivity	Human			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT. Staining of formalin-fixed tissues requires			
	boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user			
Positive control	boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal			
Positive control Shipping condition	boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-BR3 Cells. Adrenal, Lymph Node, Bladder, Prostate or			
	boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-BR3 Cells. Adrenal, Lymph Node, Bladder, Prostate or Endometrium.			
Shipping condition	boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-BR3 Cells. Adrenal, Lymph Node, Bladder, Prostate or Endometrium.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Shipping condition Storage Conditions	boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-BR3 Cells. Adrenal, Lymph Node, Bladder, Prostate or Endometrium.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when			
Shipping condition Storage Conditions Shelf life	boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-BR3 Cells. Adrenal, Lymph Node, Bladder, Prostate or Endometrium.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user SK-BR3 Cells. Adrenal, Lymph Node, Bladder, Prostate or Endometrium.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 9.8SA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified BSA-free: 1 mg/mL in PBS/0.05% azide, Purified BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified BSA-free: 1 mg/mL in PBS/0.05% azide, Purifi			

Chinese hamster ovary cells. Inquire for the specific lot.

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on December 22, 2025 at 06:42:44 AM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/interleukin-10-recombinant-monoclonal-rabbit-antibody-il10-2651r/">https://biotium.com/product/interleukin-10-recombinant-monoclonal-rabbit-antibody-il10-2651r/</a>