## **Estrogen Receptor, Alpha Recombinant** Monoclonal Mouse Antibody (rESR1/1935)



## **Product Description**

This MAb is specific to ER alpha and shows minimal cross-reaction with other members of the family. ER is an important regulator of growth and differentiation in the mammary gland. Presence of ER in breast tumors indicates an increased likelihood of response to anti-estrogen (e.g. tamoxifen) therapy. This MAb is excellent for staining of formalin-fixed, paraffin-embedded breast carcinomas. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 2300, Anti-Estrogen Receptor, alpha (rESR1/1935)

Call us: 800-304-5357

Antibody number	#2300		
Antibody reactivity	Estrogen Receptor, alpha		
(target) Antibody type	Primary		
Host species	Mouse		
Clonality	Recombinant Monoclonal		
Clone	rESR1/1935		
Isotype	IgG1		
Molecular weight	67 kDa		
Synonyms	Estrogen Receptor alpha delta 4*5,6,7*/654 isoform; Estrogen Recepto alpha delta 4 +49 isoform; Nuclear receptor subfamily 3 group A member 1		
Human gene symbol	ESR1 2099		
Entrez gene ID			
SwissProt	P03372		
Unigene	208124		
Immunogen	Recombinant full-length human Estroger Receptor alpha protein		
Antibody target cellular	Nucleus		
Verified antibody applications Species reactivity	IHC (FFPE) (verified), WB (verified)		
	Human		
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 1-2 ug/mL, Optimal dilution for a specific application should be determined by user		
Positive control	MCF-7 Cells or Breast Cancers.		
Shipping condition	Room temperature		
Storage Conditions	Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C		
Shelf life	Guaranteed for at least 24 months from date of receipt when stored as recommended		
Regulatory status	For research use only (RUO)		
Antibody/conjugate formulation	Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide Purified, BSA-free: 1 mg/mL in PBS without azide		
Antibody research areas	Cancer, Endocrinology, Nuclear receptors		
Tumor expression	Breast cancer		

Email: btinfo@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	<b>Detection channel</b>	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified.	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.

BSA-free