TACSTD2 / TROP2 Monoclonal Mouse Antibody (TACSTD2/2152)



Product Description

TACSTD2 is a cell surface glycoprotein receptor. It is a single pass type I membrane protein containing one thyroglobulin type-1 domain, an epidermal growth factor-like repeat, a phosphatidylinositol binding site and tyrosine phosphorylation sites near the C-terminus. It plays a role in transducing intracellular calcium signals. It is expressed in trophoblast cells, cornea and multi-stratified epithelia. It is also highly expressed in several types of tumors and is involved in regulating the growth of carcinoma cells.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 2152, Anti-TACSTD2 / TROP2 (TACSTD2/2152)

Product attributes

Product attributes			
Antibody number	#2152		
Antibody reactivity (target)	TACSTD2, TROP2		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	TACSTD2/2152		
Isotype	lgG1, kappa		
Molecular weight	40 kDa		
Synonyms	Cell surface glycoprotein Trop-2; Membrane Component Chromosome 1, Surface Marker 1 (M151); Pancreatic Carcinoma Marker Protein GA733-1; TROP2; Tumor-Associated Calcium Signal Transducer 2 (TACSTD2)		
Human gene symbol	TACSTD2		
Entrez gene ID	4070		
SwissProt	P09758		
Unigene	23582		
Immunogen	Recombinant fragment of human TACSTD2 protein (around aa 31-274) (exact sequence is proprietary)		
Verified antibody applications	IF (verified), IHC (FFPE) (verified)		
Antibody target cellular localization	Plasma membrane		
Species reactivity	Human		
Species reactivity Antibody application notes	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT. Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user		
	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application		
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user		
Antibody application notes Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user		
Antibody application notes Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HT29 cells. Breast or Colon Carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
Antibody application notes Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes (blowed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HT29 cells. Breast or Colon Carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when		
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HT29 cells. Breast or Colon Carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended		
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HT29 cells. Breast or Colon Carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Pmg/mL in PBS/0.05% BSA/0.05% BSA/0.05% azide, Pmg/mL in PBS/0.05% BSA/0.05% BSA/0.05% BSA/0.05% BSA/0.05% BSA/0.0		
Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HT29 cells. Breast or Colon Carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/, Purified: 0.2 mg/mL in PBS/0.05% BSA/, Purified: 0.2 mg/mL in PBS/0.05% BSA/.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% azide, Purified: 0.2 mg/mL		
Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Validated in protein array	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunohistology (formalin): 1-2 ug/mL for 30 minutes at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user HT29 cells. Breast or Colon Carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS without azide Monospecific		

Call us: 800-304-5357 Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on August 26, 2025 at 04:21:41 PM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/tacstd2-trop2-monoclonal-mouse-antibody-tacstd2-2152/