Cytokeratin, Pan Monoclonal Mouse Antibody (PCK/3150)



Product Description

Anti-cytokeratin clone PCK/3150 demonstrates a broad spectrum of cytokeratin reactivity. In normal tissues, PCK/3150 is reactive with most epithelial types, including bile ducts and hepatocytes in liver, bladder epithelium, breast ducts, bronchial epithelium, endometrium, intestinal epithelium of stomach, duodenum, ileum, colon, rectum, pancreas, ovarian epithelium, pancreatic acini, pituitary acini, pneumocytes, prostate, thyroid, skin (positive on the basal layer and negative on the superficial layers of squamous epithelium), and apocrine and sweat glands. In tumors, PCK/3150 is reactive with most carcinomas, including breast, transitional cell (TCC), renal cell (RCC), lung adenocarcinoma, lung small cell, lung squamous cell, endometrial, prostate, ovarian, hepatocellular (HCC), colorectal CA, stomach and thyroid. It is negative in certain normal tissues, including brain, lymphocytes and all cells of hematolymphoid origin, muscle, brain, nerves, endothelium and in certain tumors including most melanomas, sarcomas, lymphomas, primitive neuroectodermal tumors (PNET)/Ewings and gastrointestinal stromal tumors (GIST). Positivity has been seen on some dendritic cells in lymph nodes, some endothelia, and some muscle cells. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before p

Product attribut Antibody number

Call us: 800-304-5357

| Product attributes | | | | |
|---|---|--|--|--|
| Antibody number | #3150 | | | |
| Antibody reactivity | Cytokeratin, pan | | | |
| (target) Antibody type | Primary | | | |
| Host species | Mouse | | | |
| Clonality | Monoclonal | | | |
| Clone | PCK/3150 | | | |
| Isotype | IgG2a, kappa | | | |
| Molecular weight | 40-67 kDa (Multiple) | | | |
| Synonyms | K1B; KRT1B; Keratin; type II cytoskeletal 1b; K77; CK-1B; Keratin 1B; Keratin-77; Cytokeratin-1B; Type-II Keratin Kb39 | | | |
| Human gene symbol | KRT | | | |
| Entrez gene ID | Multiple | | | |
| SwissProt | Multiple | | | |
| Unigene | Multiple | | | |
| Immunogen | Crude cytokeratin extract prepared from RT-4 and MCF-7 cells. | | | |
| Antibody target cellular | Cytoskeleton Flow (intracellular) (verified), IF (verified), IHC (FFPE) (verified) | | | |
| Verified antibody applications | | | | |
| Species reactivity | Cat, Dog, Human | | | |
| | | | | |
| Antibody application notes | For coating for ELISA, order Ab without BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry | | | |
| Antibody application notes Positive control | BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml. for most applications, or 1 ug/million cells/100 u.L | | | |
| notes | BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml. for most applications, or 1 ug/million cells/100 uL for flow cytometry Skin, adeno- or squamous carcinomas | | | |
| Positive control | BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml. for most applications, or 1 ug/million cells/100 uL for flow cytometry Skin, adeno- or squamous carcinomas (IHC). | | | |
| Positive control Shipping condition | BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml. for most applications, or 1 ug/million cells/100 uL for flow cytometry Skin, adeno- or squamous carcinomas (IHC). Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store | | | |
| Positive control Shipping condition Storage Conditions | BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml. for most applications, or 1 ug/million cells/100 uL for flow cytometry Skin, adeno- or squamous carcinomas (IHC). Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C | | | |
| Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate | BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection using primary antibody. Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml. for most applications, or 1 ug/million cells/100 uL for flow cytometry Skin, adeno- or squamous carcinomas (IHC). Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA-free: 1 mg/mL in PBS/w. azide, Purified, BSA-free: 1 mg/mL in PBS/w. azide Guaranteed for at least 24 months from | | | |
| Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation | BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml for most applications, or 1 ug/million cells/100 uL for flow cytometry Skin, adeno- or squamous carcinomas (IHC). Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purffied: 0.2 mg/mL in PBS/0.05% BSA, Purffied: 0.2 mg/mL in PBS/0.05% BSA, Free: 1 mg/mL in PBS without azide | | | |
| Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation | BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml. for most applications, or 1 ug/million cells/100 uL for flow cytometry Skin, adeno- or squamous carcinomas (IHC). Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purffied: 0.2 mg/mL in PBS/0.05% BSA, Purffied: 0.2 mg/mL in PBS/0.05% BSA, Purffied: 0.2 mg/mL in PBS/0.5% BSA/0.05% azide, PURFIGURENT of the processor of the purffied shafes from the past 24 months from date of receipt when stored as | | | |

Email: btinfo@biotium.com

| Antibody # prefix | Conjugation | Ex/Em (nm) | Laser line | Detection channel | Dye Features |
|-------------------|-------------|------------|------------|-----------------------------|------------------|
| BNC04 | CF®405S | 404/431 | 405 | DAPI (microscopy), AF405 | CF®405S Features |
| BNC88 | CF®488A | 490/515 | 488 | GFP, FITC | CF®488A Features |
| BNC68 | CF®568 | 562/583 | 532, 561 | RFP, TRITC | CF®568 Features |
| BNC94 | CF®594 | 593/614 | 561 | Texas Red® | CF®594 Features |
| BNC40 | CF®640R | 642/662 | 633-640 | Cy®5 | CF®640R Features |
| BNC47 | CF®647 | 650/665 | 633-640 | Cy®5 | CF®647 Features |
| BNCB | Biotin | N/A | N/A | N/A | |
| BNUB | Purified | N/A | N/A | N/A | |
| BNUM | Purified, | N/A | N/A | N/A | |

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.

BSA-free