TGF-Beta Monoclonal Mouse Antibody (TGFB/510)

Product Description

This MAb recognizes TGF beta 1, 2 and 3. TGFs inhibit the growth of epithelial cells and stimulate the growth of mesenchymal cells. Most breast lesions, benign and malignant, involve abnormal proliferation and altered architecture of stromal and/or epithelial elements. Catalog number key for antibody number 0510, Anti-TGF-Beta (TGFB/510)

Call us: 800-304-5357 Email: techsupport@biotium.com

Product attributes

Product attributes				
Antibody number	#0510			
Antibody reactivity (target)	TGF-Beta			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	TGFB/510			
Isotype	lgG1, kappa			
Molecular weight	~13 kDa			
Synonyms	ARVD; BSC 1 cell growth inhibitor; CED; Cetermin; Diaphyseal dysplasia 1 (DPD1); Cliobhastoma derived T cell suppressor factor (G TSF); Latency-associated peptide (LAP); Polyergin; Transforming growth factor beta 1 (TGF beta1 or TGFB1); TGF beta2 (TGFB2); TGF beta3 (TGFB3); TGFbeta (TGFB)			
Human gene symbol	TFGB			
Entrez gene ID	7040; 7042; 7043			
SwissProt	P01137; P10600; P61812			
Unigene	645227			
Immunogen	Full length recombinant human TGF-beta2 protein			
Antibody target cellular localization	Secreted (extracellular)			
Species reactivity	Cow, Dog, Hamster, Human, Monkey, Mouse, Rat			
Antibody application notes	antibody conjugates than for indirect detection with secondary			
, ,,,	concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most			
Positive control	concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most			
	concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry Ovarian carcinoma, lung tumor or skin. PC-3 cell line. Eahy926			
Positive control	concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/ml. for most applications, or 1 ug/million cells/100 uL. for flow cytometry Ovarian carcinoma, lung tumor or skin. PC-3 cell line. Eahy926 cells.			
Positive control Shipping condition	concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry Ovarian carcinoma, lung tumor or skin. PC-3 cell line. Eahy926 cells. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Positive control Shipping condition Storage Conditions	concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry Ovarian carcinoma, lung tumor or skin. PC-3 cell line. Eahy926 cells. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C			
Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate	concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry Ovarian carcinoma, lung tumor or skin. PC-3 cell line. Eahy926 cells. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; DSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; DSA-free: 1 mg/mL in PBS/0.05% azide, Purified; DSA-free: 1 mg/mL in PBS/0.05% azide, Purified; DSA-free: 1			
Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation	concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry Ovarian carcinoma, lung tumor or skin. PC-3 cell line. Eahy926 cells. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/ree: 1 mg/mL in PBS without azide Guaranteed for at least 24 months from date of receipt when			
Positive control Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation Shelf life	concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry Ovarian carcinoma, lung tumor or skin. PC-3 cell line. Eahy926 cells. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified BSA-free: 1 mg/mL in PBS without azide Guaranteed for at least 24 months from date of receipt when stored as recommended Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on August 28, 2025 at 04:26:32 PM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/tgf-beta-monoclonal-mouse-antibody-tgfb-510/