

Prostate Specific Antigen / PSA Monoclonal Mouse Antibody (KLK3/801 + KLK3/1248)

Product Description

Recognizes a single protein of 33-34 kDa, identified as the prostate specific antigen (PSA). This MAb is highly specific to PSA and stains prostatic secretory and ductal epithelium in both normal and neoplastic tissues. PSA is a chymotrypsin-like serine protease (kallikrein family) exclusively produced by the prostate epithelium, and abundant in seminal fluid. PSA can be detected in the sera of patients with prostatic carcinoma. It is predominantly complexed to a liver-derived serine protease inhibitor, alpha-1-antichymotrypsin (ACT). A higher proportion of serum PSA is complexed to ACT in prostate cancer than in benign prostate hyperplasia. **Catalog number key for antibody number 1256, Anti-Prostate Specific Antigen (KLK3/801 KLK3/1248)**

Call us : 800-304-5357 Email:

Email: techsupport@biotium.com

Product attributes

Antibody number	#1256		
Antibody reactivity (target)	Prostate Specific Antigen		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	KLK3/801 + KLK3/1248		
Isotype	IgG1, kappa + IgG1, kappa		
Molecular weight	33-34 kDa		
Synonyms	Antigen, prostate-specific (APS), Gamma-seminoprotein, hK3, Kallikrein related peptidase 3, Kallikrein-3 (KLK3), KLK2A1, P-30 antigen, Semenogelase, Seminin		
Human gene symbol	KLK3		
Entrez gene ID	354		
SwissProt	P07288		
Unigene	171995		
Immunogen	Recombinant full-length human KLK3 protein		
Antibody target cellular localization	Secreted (extracellular)		
Verified antibody applications	IHC (FFPE) (verified)		
Species reactivity	Human		
Antibody application notes	Higher concentration may be required for direct detection using		
	primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user		
Positive control	secondary antibody, İmmunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined		
Positive control Shipping condition	secondary antibody, İmmunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user		
	secondary antibody, İmmunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user PC12 cells. Normal prostate or prostate carcinoma.		
Shipping condition	secondary antibody, İmmunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate butfer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user PC12 cells. Normal prostate or prostate carcinoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
Shipping condition Storage Conditions	secondary antibody, İmmunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user PC12 cells. Normal prostate or prostate carcinoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C		
Shipping condition Storage Conditions Regulatory status Antibody/conjugate	secondary antibody, İmmunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed itsues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user PC12 cells. Normal prostate or prostate carcinoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in		
Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation	secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user PC12 cells. Normal prostate or prostate carcinoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide		
Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation Shelf life	secondary antibody, Immunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user PC12 cells. Normal prostate or prostate carcinoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Guaranteed for at least 24 months from date of receipt when stored as recommended		
Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation Shelf life Cell/tissue expression	secondary antibody, İmmunohistochemistry (formalin-fixed): 0.5-1.0 ug/mL for 30 min at RT, Staining of formalin-fixed itsues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user PC12 cells. Normal prostate or prostate carcinoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS without azide Guaranteed for at least 24 months from date of receipt when stored as recommended Prostate Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in		

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on September 17, 2025 at 08:11:10 AM. Visit product page to check for updated information before use. Product link: <u>https://biotium.com/product/prostate-specific-antigen-psa-monoclonal-mouse-antibody-klk3-801-klk3-1248/</u>