## p57 / KIP2 Monoclonal Mouse Antibody (57P06)

## **Product Description**

Recognizes a protein of 57 kDa, identified as p57Kip2. It shows no cross-reaction with p27Kip1. p57Kip2 is a potent tight-binding inhibitor of several G1 cyclin complexes, and is a negative regulator of cell proliferation. Anti-p57 has been used as an aide in identification of complete hydatidiform mole (CHM) (no nuclear labeling of cytotrophoblasts and stromal cells) from partial hydatidiform mole (PHM) in which both cytotrophoblasts and stromal cells stain. The histological differentiation of complete mole, partial mole, and hydropic spontaneous abortion is problematic. Most complete hydatidiform moles are diploid, whereas most partial moles are triploid. Ploidy studies will identify partial moles, but will not differentiate complete moles from non-molar gestations. Complete moles carry a high risk of persistent disease and choriocarcinoma, while partial moles have a very low risk. In normal placenta, many cytotrophoblast nuclei and stromal cells are labeled with this antibody. Similar findings apply to PHM and hydropic abortus tissues. Intervillous trophoblastic islands (IVTIs) demonstrate nuclear labeling in all three entities and serve as an internal control. Catalog number key for antibody number 1264, Anti-KIP2|p57 (57P06)

## Call us: 800-304-5357 Email: techsupport@biotium.com

## **Product attributes**

Antibody number	#1264			
Antibody reactivity (target)	KIP2, p57			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	57P06			
Isotype	IgG2b, kappa			
Molecular weight	57 kDa			
Synonyms	Beckwith Wiedemann syndrome (WBS); BWCR; Cyclin dependent kinase inhibitor 1C (CDKN1C); KIP2; p57			
Human gene symbol	CDKN1C			
Entrez gene ID	1028			
SwissProt	P49918			
Unigene	106070			
Immunogen	Recombinant full-length human p57Kip2 protein			
Verified antibody applications	IHC (FFPE) (verified)			
Antibody target cellular localization	Nucleus			
Species reactivity	Human. Mouse.			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistochemistry (formalin-fixed): 0.25-0.5 ug/mL for 30 minutes at RT, Flow cytometry: 0.5-1 ug/million cells, immunofluorescence: 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 minutes followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user			
Danishing and and				
Positive control	LS174T, Raji, HT29, SK-BR3 cells. Colon or Prostate carcinomas.			
Shipping condition				
	carcinomas.			
Shipping condition	carcinomas. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Shipping condition Storage Conditions	carcinomas. Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C			
Shipping condition Storage Conditions Regulatory status Antibody/conjugate	carcinomas. Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purifiled: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purifiled, BSA-free: 1 mg/mL in PBS/0.05%			
Shipping condition Storage Conditions Regulatory status Antibody/conjugate formulation	carcinomas.  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide  Guaranteed for at least 24 months from date of receipt when			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on August 5, 2025 at 01:50:57 AM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/p57-kip2-monoclonal-mouse-antibody-57p06/">https://biotium.com/product/p57-kip2-monoclonal-mouse-antibody-57p06/</a>