## CD99 Monoclonal Mouse Antibody (HO36-1.1)



## **Product Description**

Recognizes a sialoglycoprotein of 27-32 kDa, identified as CD99, or MIC2 gene product, or E2 antigen. MIC2 gene is located in the pseudo-autosomal region of the human X and Y chromosome. MIC2 gene encodes two distinct proteins, which are produced by alternative splicing of the CD99 gene transcript and are identified as bands of 30 and 32 kDa (p30/32). Although its function is not fully understood, CD99 is implicated in various cellular processes including homotypic aggregation of T cells, upregulation of T cell receptor and MHS molecules, apoptosis of immature thymocytes and leukocyte diapedesis.CD99 is expressed on the cell membrane of some lymphocytes, cortical thymocytes, and granulosa cells of the ovary. Most pancreatic islet cells, Sertoli cells of the testis, and some endothelial cells express this antigen. Mature granulocytes express very little or no CD99. MIC2 is strongly expressed on Ewing's sarcoma cells and primitive peripheral neuroectodermal tumors. This MAb shows a very similar reactivity to other CD99 MAbs (e.g. O13, 12E7, or HBA-71) and is excellent for immunohistochemical staining of formalin-fixed, paraffin-embedded tissues.

This antibody is available purified with BSA/azide at 200 ug/mL, or BSA/azide-free at 1 mg/mL.

Catalog number key for antibody number 0869, Anti-CD99 (HO36-1.1)

## Call us : 800-304-5357

Product attributes				
Antibody number	#0869			
Antibody reactivity (target)	CD99			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	HO36-1.1			
Isotype	IgM, kappa			
Molecular weight	27-32 kDa			
Synonyms	12E7; E2 antigen; MIC 2X; MIC 2Y; MIC2; Protein MIC2; Surface antigen MIC2; T-cell surface glycoprotein E2			
Human gene symbol	CD99			
Entrez gene ID	4267			
SwissProt	P14209			
Unigene	653349			
Immunogen	Purified E-rosette forming cells from human peripheral blood lymphocytes			
Verified antibody applications	IHC (FFPE) (verified)			
Species reactivity	Human, Rat			
Antibody target cellular localization	Plasma membrane			
Antibody application notes	Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user			
Antibody application notes  Shipping condition	formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific			
	formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user			
Shipping condition	formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  Room temperature  Store at 2 to 8 °C, Note: store BSA-free antibodies at -10 to -35			
Shipping condition Storage Conditions	formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  Room temperature  Store at 2 to 8 °C, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when			
Shipping condition Storage Conditions Shelf life	formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  Room temperature  Store at 2 to 8 °C, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended			
Shipping condition Storage Conditions Shelf life Regulatory status	formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  Room temperature  Store at 2 to 8 °C, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody research areas Antibody/conjugate	formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  Room temperature  Store at 2 to 8 °C, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Cancer, Endocrinology, Immunology  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, PIRFe: 1 mg/mL in			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody research areas Antibody/conjugate formulation	formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  Room temperature  Store at 2 to 8 °C, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Cancer, Endocrinology, Immunology  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody research areas Antibody/conjugate formulation Positive control	formalin-fixed 1-2 ug/mL, Flow Cytometry 5-10 uL/million cells in 0.1 mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Optimal dilution for a specific application should be determined by user  Room temperature  Store at 2 to 8 °C, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Cancer, Endocrinology, Immunology  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide  Pancreas or Ewing's sarcoma  Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on December 12, 2025 at 07:26:33 AM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/monoclonal-anti-cd99-ho36-1-1/">https://biotium.com/product/monoclonal-anti-cd99-ho36-1-1/</a>