## Mucin 5AC / MUC 5AC Monoclonal Mouse Antibody (CLH2)



## **Product Description**

Mucin 5AC glycoprotein (MUC5AC) is a HMW glycoprotein belonging to the superfamily of mucins. Mucins are produced by epithelial cells and can be divided into two families; secretory mucins and membrane bound mucins. MUC5AC is a mucus-forming, secreted mucin that is found in normal gastric and tracheo-bronchial mucosa, but absent from normal colon. MUC5AC expression is present in primary ovarian mucinous cancer but usually absent in colorectal adenocarcinoma, thus showing an expression pattern opposite to MUC2. Together with a panel of antibodies, Anti-MUC5AC may be useful for differential identification of primary mucinous ovarian tumors from colon adenocarcinoma metastatic to the ovary. MUC5AC antibodies may also be useful for identification of intestinal metaplasia as well as in the identification of pancreatic carcinoma and pre-cancerous changes vs. normal pancreas.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0897, Anti-Mucin 5AC (CLH2)

## Product attributes

| Product attributes  |  |  |  |
|---|--|--|--|
| Antibody number   | #0897  |  |  |
| Antibody reactivity (target)  | Mucin 5AC  |  |  |
| Antibody type   | Primary  |  |  |
| Host species  | Mouse  |  |  |
| Clonality   | Monoclonal   |  |  |
| Clone   | CLH2   |  |  |
| Isotype   | IgG1, kappa  |  |  |
| Molecular weight  | >1,000 kDa   |  |  |
| Synonyms  | Apomucin Major Airway Glycoprotein, Mucin 5 subtype AC tracheobronchial, Mucin 5 Subtypes A And C, Mucin 5AC oligomeric mucus/gel forming, Tracheobronchial Mucin (TBM)  |  |  |
| Human gene symbol   | MUC5AC   |  |  |
| Entrez gene ID  | 4586   |  |  |
| SwissProt   | P98088   |  |  |
| Unigene   | 534332   |  |  |
| Immunogen   | A synthetic peptide of human MUC5AC tandem repeat  |  |  |
| Verified antibody applications  | IHC (FFPE) (verified)  |  |  |
| Antibody target cellular localization   | Secreted (extracellular)   |  |  |
|   |  |  |  |
| Species reactivity  | Human  |  |  |
| Species reactivity  Antibody application notes  | Human  Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optima (Idution for a specific application should be determined by user  |  |  |
| •   | Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific   |  |  |
| Antibody application notes  | Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user  |  |  |
| Antibody application notes  Positive control  | Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mt, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user  |  |  |
| Antibody application notes  Positive control Shipping condition   | Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user  Stomach  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,   |  |  |
| Antibody application notes  Positive control Shipping condition Storage Conditions  | Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RTI for 20 minutes, Flow Cytometry 0.5-1 ug/milion cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user  Stomach  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when   |  |  |
| Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life   | Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/milion cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user  Stomach  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  |  |  |
| Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate                                      | Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/milion cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user  Stomach  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.5 mg/mL i |  |  |
| Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation                          | Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RTf for 20 minutes, Flow Cytometry 0.5-1 ug/milion cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user  Stomach  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide   |  |  |
| Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation  Antibody research areas | Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/mlilion cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user  Stomach  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide  Cancer, Mucins  Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in   |  |  |

Call us: 800-304-5357 Email: techsupport@biotium.com

| Antibody # prefix | Conjugation           | Ex/Em (nm) | Laser line | Detection channel           | Dye Features     |
|-------------------|-----------------------|------------|------------|-----------------------------|------------------|
| BNC04             | CF®405S               | 404/431    | 405        | DAPI (microscopy),<br>AF405 | CF®405S Features |
| BNC88             | CF®488A               | 490/515    | 488        | GFP, FITC                   | CF®488A Features |
| BNC68             | CF®568                | 562/583    | 532, 561   | RFP, TRITC                  | CF®568 Features  |
| BNC94             | CF®594                | 593/614    | 561        | Texas Red®                  | CF®594 Features  |
| BNC40             | CF®640R               | 642/662    | 633-640    | Cy®5                        | CF®640R Features |
| BNC47             | CF®647                | 650/665    | 633-640    | Cy®5                        | CF®647 Features  |
| BNC74             | CF®740                | 742/767    | 633-685    | 775/50                      | CF®740 Features  |
| BNCB              | Biotin                | N/A        | N/A        | N/A                         |                  |
| BNUB              | Purified              | N/A        | N/A        | N/A                         |                  |
| BNUM              | Purified,<br>BSA-free | N/A        | N/A        | N/A                         |                  |

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on July 12, 2025 at 09:24:59 PM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/monoclonal-anti-mucin-5ac-muc5ac-clh2/