CD45RB Monoclonal Mouse Antibody (BRA-11G)



Product Description

CD45R, also designated CD45 and PTPRC, has been identified as a transmembrane glycoprotein, broadly expressed among hematopoietic cells. Multiple isoforms of CD45R are distributed throughout the immune system according to cell type. These isoforms arise because of alternative splicing of exons 4, 5, and 6. The corresponding protein domains are characterized by the binding of monoclonal antibodies specific for CD45RA (exon 4), CD45RB (exon 5), CD45RC (exon 6) and CD45RO (exons 4 to 6 spliced out). The variation in these isoforms is localized to the extracellular domain of CD45R, while the intracellular domain is conserved. CD45RB is expressed on mature B-lymphocytes and the majority of lymphomas and leukemias of B-cell origin.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0174, Anti-CD45RB (BRA-11G)

Product attributes

Call us: 800-304-5357

Product attributes			
Antibody number	#0174		
Antibody reactivity (target)	CD45RB		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	BRA-11G		
Isotype	IgG1, kappa		
Molecular weight	180-220 kDa		
Synonyms	B220, CD45R, GP180, Leukocyte common antigen (LCA), Loc Ly-5, Lyt-4, Protein tyrosine phosphatase receptor type C (PTPRC), Receptor-type tyrosine-protein phosphatase C, T200 glycoprotein		
Human gene symbol	PTPRC		
Entrez gene ID	5788		
SwissProt	P08575		
Unigene	654514		
Immunogen	Non-T, non-B CALLA positive ALL cell line REH (Leucocyte Workshop IV and V)		
Antibody target cellular localization	Plasma membrane		
Verified antibody applications	Flow (verified), IHC (FFPE) (verified), WB (verified)		
Species reactivity	Human		
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
Antibody application notes Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a		
	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
Positive control Shipping condition	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Ramos, U-698, or GA-10 cells. Tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
Positive control Shipping condition Storage Conditions	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Ramos, U-698, or GA-10 cells. Tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when		
Positive control Shipping condition Storage Conditions Shelf life	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Ramos, U-698, or GA-10 cells. Tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended		
Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Ramos, U-698, or GA-10 cells. Tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified in PBS/0.05% BSA/0.05% azide, Purified in PBS/0.05% BSA/0.05% azide, Purified: 1 mg/mL in PBS/0.05% BSA/0.05% azide, PIRIFICATE To Mg/mL in PBS/0.05% BS		
Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Ramos, U-698, or GA-10 cells. Tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS without azide		
Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Antibody research areas	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Does not react with rat or rhesus monkey; others not known, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user Ramos, U-698, or GA-10 cells. Tonsil Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Immunology, Signal transduction Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in		

Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on August 27, 2025 at 06:52:45 AM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/monoclonal-anti-cd45rb-bra-11g/