Myogenin Monoclonal Mouse Antibody (F5D)



Product Description

Myogenin is a member of the MyoD family of myogenic basic helix-loop-helix (bHLH) transcription factors that also includes MyoD, Myf-5, and MRF4 (also known as herculinor Myf-6). MyoD family members are expressed exclusively in skeletal muscle and play a key role in activating myogenesis by binding to enhancer sequences of muscle-specific genes. The regulatory domain of MyoD is approximately 70 amino acids in length and includes both a basic DNA binding motif and a bHLH dimerization motif. MyoD family members share about 80% amino acid homology in their bHLH motifs. Anti-myogenin labels the nuclei of myoblasts in developing muscle tissue, and is expressed in tumor cell nuclei of rhabdomyosarcoma and some leiomyosarcomas. Positive nuclear staining may occur in Wilms tumor. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. **Stock status**: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. **Catalog number key** for antibody number 0632, Anti-Myogenin (F5D)

Tumor expression

Call us: 800-304-5357

Product attributes				
Antibody number	#0632			
Antibody reactivity	Myogenin			
(target) Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	F5D			
Isotype	IgG1, kappa			
Molecular weight	34 kDa			
Synonyms	bHLHc3, cb553, Class C basic helix-loop-helix protein 3, Myf-4, MYF4, MYOG, Myogenic factor 4			
Uuman sana aymbal	MYOG			
Human gene symbol				
Entrez gene ID	4656 P45470			
SwissProt	P15173			
Unigene	2830			
Immunogen	Rat myogenin peptide (aa 73-94) followed by rat myogenin recombinant fragment (aa30-224) (Epitope aa138-158)			
Antibody target cellular	Nucleus			
Verified antibody	IHC (FFPE) (verified)			
applications Species reactivity	Cat, Dog, Human, Mouse, Pig, Rat			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunoprecipitation: 0.5-1 ug/500 ug protein lysate, Immunofluorescence: 0.5-1 ug/ml, Immunohistology (formalin), Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/mlllion cells/0.1 ml., Western blotting 0.5-1.0 ug/ml. ELISA For coating, order Ab without BSA, Optimal dilution for a specific application should be determined by user			
	without BSA, Optimal dilution for a specific application should be			
Positive control	without BSA, Optimal dilution for a specific application should be			
Positive control Shipping condition	ug/mL, ELISA For coating, order Ab without BSA, Optimal dilution for a specific application should be determined by user Rh-30 cells. Skeletal muscle or			
	ug/mL, ELISA For coating, order Ab without BSA, Optimal dillution for a specific application should be determined by user Rh-30 cells. Skeletal muscle or rhabdomyosarcoma.			
Shipping condition	ug/mL, ELISA Por coating, order Ab without BSA, Optimal dillution for a specific application should be determined by user Rh-30 cells. Skeletal muscle or rhabdomyosarcoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as			
Shipping condition Storage Conditions Shelf life	ug/mL, ELISA For coating, order Ab without BSA, Optimal dilution for a specific application should be determined by user Rh-30 cells. Skeletal muscle or rhabdomyosarcoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			
Shipping condition Storage Conditions	ug/mL, ELISA For coating, order Ab without BSA, Optimal dillution for a specific application should be determined by user Rh-30 cells. Skeletal muscle or rhabdomyosarcoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO)			
Shipping condition Storage Conditions Shelf life	ug/mL, ELISA For coating, order Ab without BSA, Optimal dilution for a specific application should be determined by user Rh-30 cells. Skeletal muscle or rhabdomyosarcoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	ug/mL, ELISA Por coating, order Ab without BSA, Optimal dilution for a specific application should be determined by user Rh-30 cells. Skeletal muscle or rhabdomyosarcoma. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/oree: 1 mg/mL in PBS/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS/0.16%			

Email: btinfo@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.