

CD35 / CR1 Monoclonal Mouse Antibody (E11)



Product Description

Recognizes a protein of 210-220 kDa, which is identified as the complement receptor 1 (CR1)/CD35. This MAb is specific for a site in CR1 that is distal from the C3b-binding site, so that it is unable to block CR1 activity. This MAb is highly specific to CR1 and shows no cross-reaction with CR2. The primary function of CR1 is to serve as the cellular receptor for C3b and C4b, the most important components of the complement system leading to clearance of foreign macromolecules. The Knops blood group system is a system of antigens located on this protein. Follicular dendritic cells (FDC) are restricted to the B-cell regions of secondary lymphoid follicles. They are CD21 /CD35 /CD1a-. This MAb labels follicular dendritic cells and follicular dendritic cell sarcoma.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0040, Anti-CD35 (E11)

Call us : 800-304-5357 Email: techsupport@biotium.com

Product attributes

Product attributes			
Antibody number	#0040		
Antibody reactivity (target)	CD35, CR1		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	E11		
Isotype	IgG1, kappa		
Molecular weight	210-220 kDa		
Synonyms	C3 binding protein, C3b/C4b receptor, C3BR, C4BR, Complement Component (3b/4b) receptor 1 including Knops blood group system, Complement receptor type 1, KN, Knops blood group antigen		
Human gene symbol	CR1		
Entrez gene ID	1378		
SwissProt	P17927		
Unigene	334019		
Immunogen	Intact human monocytes		
Antibody target cellular localization	Plasma membrane		
Verified antibody applications	IHC (FFPE) (verified)		
Species reactivity	Baboon, Cynomolgus monkey, Human, Rhesus monkey		
Expected antibody applications	ELISA (published for clone), Flow, surface (published for clone), WB (published for clone)		
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immonfluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
Positive control	Follicular dendritic cells (FDC) in tonsil		
Shipping condition	Room temperature		
Storage Conditions	Store at 2 to 8 $^\circ$ C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 $^\circ$ C		
Shelf life	Guaranteed for at least 24 months from date of receipt when stored as recommended		
Regulatory status	For research use only (RUO)		
Antibody/conjugate formulation	Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide		
Antibody research areas	Immunology		
Product origin	Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot.		
Cell/tissue expression	Antigen presenting cells		

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

References

Note: References for this clone sold by other suppliers may be listed for expected applications.

- 1. J Immunol (1999) 162:7549-7554. (WB) 2. Clin Exp Immunol (2005) 143: 274-280. (ELISA) 3. Int Immunol (2013) 25(1): 25-33. (Flow, surface)

This datasheet was generated on August 1, 2025 at 07:34:15 AM. Visit product page to check for updated information before use. Product link: <u>https://biotium.com/product/monoclonal-anti-cd35-cr1-e11/</u>