Cytokeratin 8 Monoclonal Mouse Antibody (KRT8/803)



Product Description

Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). CK8 is primarily found in the non-squamous epithelia and is present in majority of adenocarcinomas and ductal carcinomas. It is absent in squamous cell carcinomas. Hepatocellular carcinomas are defined by the use of antibodies that recognize only cytokeratin 8 and 18. CK8 exists on several types of normal and neoplastic epithelia, including many ductal and glandular epithelia such as colon, stomach, small intestine, trachea, and esophagus as well as in transitional epithelium. Anti-CK8 does not react with skeletal muscle or nerve cells. Epithelioid sarcoma, chordoma, and adamantinoma show strong positivity corresponding to that of simple epithelia (with antibodies against CK8, CK18 and CK19). Anti-CK8 is useful for the differentiation of lobular (ring-like, perinuclear) from ductal (peripheral-predominant) carcinoma of the breast. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. **Stock status:** Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before also incurred to Confusion and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 0803, Anti-Cytokeratin 8

Call us: 800-304-5357

| Product attributes | | | | |
|------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--|--|--|
| Antibody number | #0803 | | | |
| Antibody reactivity | Cytokeratin 8 | | | |
| (target) Antibody type | Primary | | | |
| Host species | Mouse | | | |
| Clonality | Monoclonal | | | |
| Clone | KRT8/803 | | | |
| Isotype | IgG1, kappa | | | |
| Molecular weight | 52.5 kDa | | | |
| Synonyms | CARD2; CK8; CYK8; CYKER; Cytokeratin Endo A; DreK8; EndoA; K2C8; K8; Keratin 8; Krt 2.8; KRT8; Type-II Keratin Kb8 | | | |
| Human gene symbol | KRT8 | | | |
| Entrez gene ID | 3856 | | | |
| SwissProt | P05787 | | | |
| Unigene | 533782 & 708445 | | | |
| Immunogen | Recombinant human KRT8 protein | | | |
| Antibody target cellular | Cytoskeleton | | | |
| Verified antibody | IHC (FFPE) (verified), WB (verified) | | | |
| applications Species reactivity | Human | | | |
| Antibody application notes | Higher concentration may be required direct detection using primary antibody conjugates than for indirect detection with secondary antibody, immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1.C ug/mL, Staining of formalin-fixed dissue requires boiling tissue sections in 10 m citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minute Flow Cytometry 0.5-1 ug/million cells/0 mL, Optimal dilution for a specific application should be determined by user | | | |
| Positive control | MCF-7 or A431 cells. Skin, Colon, lung or breast carcinoma | | | |
| Shipping condition | Room temperature | | | |
| Storage Conditions | Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C | | | |
| Shelf life | Guaranteed for at least 24 months from date of receipt when stored as recommended | | | |
| Regulatory status | For research use only (RUO) | | | |
| Antibody/conjugate formulation | Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide Purified, BSA-free: 1 mg/mL in PBS without azide | | | |
| Antibody research areas | Cancer, Cytoskeleton | | | |
| Cell/tissue expression | Epithelial cells | | | |
| Tumor expression | Breast cancer, Hepatocellular carcinoma, Sarcoma | | | |

Email: btinfo@biotium.com

| Antibody # prefix | Conjugation | Ex/Em (nm) | Laser line | Detection channel | Dye Features |
|-------------------|-----------------------|------------|------------|-----------------------------|------------------|
| BNC04 | CF®405S | 404/431 | 405 | DAPI (microscopy), AF405 | CF®405S Features |
| BNC88 | CF®488A | 490/515 | 488 | GFP, FITC | CF®488A Features |
| BNC68 | CF®568 | 562/583 | 532, 561 | RFP, TRITC | CF®568 Features |
| BNC94 | CF®594 | 593/614 | 561 | Texas Red® | CF®594 Features |
| BNC40 | CF®640R | 642/662 | 633-640 | Cy®5 | CF®640R Features |
| BNC47 | CF®647 | 650/665 | 633-640 | Cy®5 | CF®647 Features |
| BNCB | Biotin | N/A | N/A | N/A | |
| BNUB | Purified | N/A | N/A | N/A | |
| BNUM | Purified, BSA-free | N/A | N/A | N/A | |

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.