## EGFR Monoclonal Mouse Antibody (B1D8)

## **Product Description**

This antibody reacts with the extracellular domain of EGFR and blocks the EGF/TGF induced activation. It also blocks tumor growth in vivo. It is excellent for purification of EGFR. EGFR is type I receptor tyrosine kinase with sequence homology to erbB-1, -2, -3 -4 or HER-1, -2, -3 -4. It binds to Epidermal Growth Factor (EGF), Transforming Growth Factor-a (TGF-a), Heparin-binding EGF (HB-EGF), amphiregulin, βcellulin and epiregulin.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

**Stock status:** Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0643, Anti-EGFR (B1D8)

## Draduat attributas

Product attributes				
Antibody number	#0643			
Antibody reactivity (target)	EGFR			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	B1D8			
Isotype	IgG2a, kappa			
Molecular weight	~170 kDa (wild type) and ~145 kDa (vIII variant)			
Synonyms	Erbb1; ERBB1; Errp; HER1; mENA; PIG61; Proto-oncogene c-ErbB-1; Receptor Tyrosine Protein Kinase; ErbB1; Urogastrone; wa2; Wa5			
Human gene symbol	EGFR			
Entrez gene ID	1956			
SwissProt	P00533			
Unigene	488293			
Immunogen	Microsomes from A431 cells.			
Antibody target cellular localization	Plasma membrane			
Species reactivity	Human			
Expected antibody applications	Flow, surface (published for clone), Functional studies (published for clone), IP (published for clone)			
	primary antibody conjugates than for indirect detection with			
Antibody application notes	secondary antibody, Does not react with mouse or rat, others not known, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Affinity Purification, for coupling, order Ab without BSA, Immunoprecipitation 1-2 ug/500 ug protein lysate, Optimal			
Antibody application notes  Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Does not react with mouse or rat, others not known, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Affinity Purification, for coupling, order Ab without BSA, Immunoprecipitation 1-2 ug/500 ug protein lysate, Optimal			
	primary antibody conjugates than for indirect detection with secondary antibody, Does not react with mouse or rat, others not known, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Affinity Purification, for coupling, order Ab without BSA, Immunoprecipitation 1-2 ug/500 ug protein lysate, Optimal dilution for a specific application should be determined by user			
Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Does not react with mouse or rat, others not known, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Affinity Purfication, for coupling, order Ab without BSA, Immunoprecipitation 1-2 ug/500 ug protein lysate, Optimal dilution for a specific application should be determined by user A431 cells, Breast or Bladder cancer			
Positive control Shipping condition	primary antibody conjugates than for indirect detection with secondary antibody, Does not react with mouse or rat, others not known, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Affinity Purification, for coupling, order Ab without BSA, Immunoprecipitation 1-2 ug/500 ug protein lysate, Optimal dilution for a specific application should be determined by user A431 cells, Breast or Bladder cancer Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Positive control Shipping condition Storage Conditions	primary antibody conjugates than for indirect detection with secondary antibody, Does not react with mouse or rat, others not known, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Affinity Purification, for coupling, order Ab without BSA, Immunoprecipitation 1-2 ug/500 ug protein lysate, Optimal dilution for a specific application should be determined by user A431 cells, Breast or Bladder cancer Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when			
Positive control Shipping condition Storage Conditions Shelf life	primary antibody conjugates than for indirect detection with secondary antibody, Does not react with mouse or rat, others not known, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Affinity Purification, for coupling, order Ab without BSA, Immunoprecipitation 1-2 ug/500 ug protein lysate, Optimal dilution for a specific application should be determined by user A431 cells, Breast or Bladder cancer Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			
Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	primary antibody conjugates than for indirect detection with secondary antibody, Does not react with mouse or rat, others not known, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Affinity Purification, for coupling, order Ab without BSA, Immunoprecipitation 1-2 ug/500 ug protein lysate, Optimal dilution for a specific application should be determined by user A431 cells, Breast or Bladder cancer Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg			
Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	primary antibody conjugates than for indirect detection with secondary antibody, Does not react with mouse or rat, others not known, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Affinity Purification, for coupling, order Ab without BSA, Immunoprecipitation 1-2 ug/500 ug protein lysate, Optimal dilution for a specific application should be determined by user A431 cells, Breast or Bladder cancer Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS without azide			

Call us: 800-304-5357 Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

## References

Note: References for this clone sold by other suppliers may be listed for expected applications.

- 1. J Biol Chem (1984) 259(19): 11895-11900. (IP, functional studies)
- 2. J Immunol Meth (2004) 287: 147-158. (Flow; epitope mapping)