## CD43 Monoclonal Mouse Antibody (84-3C1)



## **Product Description**

This antibody recognizes a cell surface glycoprotein of 95/115/135 kDa (depending upon the extent of glycosylation), identified as CD43 (Workshop III). 70-90% of T-cell lymphomas and from 22-37% of B-cell lymphomas express CD43. No reactivity has been observed with reactive B-cells. So a B-lineage population that co-expresses CD43 is highly likely to be a malignant lymphoma, especially a low-grade lymphoma, rather than a reactive B-cell population. When CD43 antibody is used in combination with anti-CD20, effective immunophenotyping of the lymphomas in formalin-fixed tissues can be obtained. Co-staining of a lymphoid infiltrate with anti-CD20 and anti-CD43 argues against a reactive process and favors a diagnosis of lymphoma.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0342, Anti-CD43 (84-3C1)

## Droduct attributes

Call us: 800-304-5357

Product attributes			
Antibody number	#0342		
Antibody reactivity (target)	CD43		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	84-3C1		
Isotype	IgG1, kappa		
Molecular weight	95, 115, or 135 kDa		
Synonyms	Galactoglycoprotein, GALGP, GPL115, Leukocyte sialoglycoprotein, Leukosialin, LSN, Sialophorin, SPN		
Human gene symbol	SPN		
Entrez gene ID	6693		
SwissProt	P16150		
Unigene	632188		
Immunogen	Stimulated human leukocytes		
Antibody target cellular localization	Plasma membrane		
Verified antibody applications	Flow (surface) (verified), IF (verified), IHC (FFPE) (verified), WB (verified)		
Species reactivity	Human		
Expected antibody applications	IP (published for clone)		
applications			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined		
Antibody application notes	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
Antibody application notes  Positive control	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  Paracortex in a tonsil or a reactive lymph node		
Antibody application notes  Positive control Shipping condition	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  Paracortex in a tonsil or a reactive lymph node  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
Antibody application notes  Positive control Shipping condition Storage Conditions	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  Paracortex in a tonsil or a reactive lymph node  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when		
Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  Paracortex in a tonsil or a reactive lymph node  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended		
Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/mlilion cells/0.1 mL, Optimal dilution for a specific application should be determined by user  Paracortex in a tonsil or a reactive lymph node  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 9.8A-free: 1 mg/mL in PBS/0.05% BSA, Purified: 9.8A-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Pmg/mL in PBS/0.05% azide, Pmg/mL in PBS/0.05% BSA/0.05% azide,		
Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/mlilion cells/0.1 mL, Optimal dilution for a specific application should be determined by user  Paracortex in a tonsil or a reactive lymph node  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS without azide		
Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation  Antibody research areas	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Optimal dilution for a specific application should be determined by user  Paracortex in a tonsil or a reactive lymph node  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% BSA/0.05% BSA/free: 1 mg/mL in PBS without azide  Immunology  Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in		
Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation  Antibody research areas Product origin	primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min tollowed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Optimal dilution for a specific application should be determined by user  Paracortex in a tonsil or a reactive lymph node  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide  Immunology  Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot.		

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

## References

Note: References for this clone sold by other suppliers may be listed for expected applications.

Eur J Immunol (1987)17: 1523-1526. (immunoprecipitation)

This datasheet was generated on December 22, 2025 at 07:05:54 PM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/monoclonal-anti-cd43-84-3c1/">https://biotium.com/product/monoclonal-anti-cd43-84-3c1/</a>