Involucrin Monoclonal Mouse Antibody (SY5)



Product Description

This antibody recognizes a protein of 66 kDa-170 kDa, identified as involucrin. In Western blotting of cultured human keratinocytes, this MAb reacts with a 120 kDa protein. It stains the involucrin in a variety of sizes: 170 kDa in MCF-7 cells, a doublet of ~115 kDa and 150 kDa in gorilla and owl monkey, 66 kDa in dog, and a doublet of 105 kDa in pig. Its epitope maps between codon 421-568 of human involucrin. Involucrin is expressed in a range of stratified squamous epithelia, including the cornea, which lacks a distinct cornified layer. In normal epidermis, it is first expressed in the upper spinous layers, and in keratinocyte cultures, all cells that have left the basal layer express it. Involucrin expression is altered in pathological conditions: in psoriasis and other benign epidermal hyperplasias, involucrin expression begins closer to the basal layer than normal; expression is abnormal in squamous cell carcinomas and premalignant lesions, and is reduced in severe dysplasias of the larynx and cervix. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 0678, Anti-Involucrin (SY5)

Call us: 800-304-5357 Email: btinfo@biotium.com

direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunohistology formalin-fixed 0.1-0.2 ug/ml., Staining of formalin-fixed 0.1-0.2 ug/ml., Staining of formalin-fixed 0.1-0.2 ug/ml., Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at 1 mg/ml. PBS for 5 min at 37C, enzyme digestion is better than citrate buffer based epitope unmasking, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Room temperature Storage Conditions Storage Conditions Storage Conditions Storage Conditions Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.18 BSA, Purified: 0.2	Product attributes				
Host species Mouse	Antibody number	#0678			
Host species Mouse Clonality Monoclonal Clone SY5 Isotype IgG1, kappa Molecular weight 66-170 kDa Synonyms IVL Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Antibody target cellular Cytoplasmic Incelliation Antibody application species reactivity Antibody application notes Antibody application on species reactive direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology formalin-fixed tissues is enhanced by digestion with trypsin or Protease XVX at Img/mL PBS for 5 min at 37C, enzyme digestion is better than citrate buffer based epitope unmasking, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions For research use only (RUO) Antibody/conjugate formulation Antibody/conjugate For research use only (RUO) Antibody/conjugates For research use only (RUO) Antibody research areas Antibody research areas Antibody research areas		Involucrin			
Clone SY5 Isotype IgG1, kappa Molecular weight 66-170 kDa Synonyms IVL Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Antibody target cellular Cytoplasmic Indication Species reactivity Antibody application notes Antibody application notes Antibody application and in a specific detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofibiology formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at Img/mL PBS for 5 min at 37C, enzyme digestion is better than citrate buffer based epitope unmasking, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofibiorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions For research use only (RUO) Antibody/conjugate for might, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Antibody research areas Antibody research areas Antibody research areas	Antibody type	Primary			
Clone SY5 Isotype IgG1, kappa Molecular weight 66-170 kDa Synonyms IVL Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Antibody target cellular Cytoplasmic Incollipation Perified antibody application species reactivity Antibody application notes Antibody application notes Antibody application application species reactivity Antibody application notes Antibody application and in the properties of the properties	Host species	Mouse			
Isotype	Clonality	Monoclonal			
Molecular weight Synonyms IVL Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Antibody target cellular Verified antibody applications Species reactivity Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at Img/mL PBS for 5 min at 37c, enzyme digestion is better than citrate buffer based epitope ummasking, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions For research use only (RUO) Antibody/conjugate formulation For research use only (RUO) Antibody/conjugate formulation Antibody research areas Antibody research areas Antibody research areas Cancer	Clone	SY5			
Synonyms IVL Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Antibody target cellular Cytoplasmic Perified antibody application Species reactivity Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunohistology formalin-fixed tissues is enhanced by digestion with trypsin or Prolease XXV at 1mg/mL PBS for 5 min at 37c, erzyme digestion is better than citrate buffer based epitope ummasking, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Antibody research areas Antibody research areas Antibody research areas	Isotype	IgG1, kappa			
Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Antibody target cellular Cytoplasmic Perified antibody application species reactivity Antibody application notes Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at Img/mL PBS for 5 min at 37C, enzyme digestion is better than citrate buffer based epitope urmasking, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions Storage Conditions For eact vith most, others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions For at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation PBS/0.05% BSA/0.05% BSA/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS	Molecular weight	66-170 kDa			
Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Antibody target cellular Cytoplasmic Purified antibody application Species reactivity Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunohistology formalin-fixed 0.1-0.2 ug/mL, Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Prolease XXV at 1mg/mL PBS for 5 min at 37c, erzyme digestion is better than citrate buffer based epitope ummasking, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Room temperature Storage Conditions Storage Conditions Storage Conditions Storage Conditions For eact vith mose; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions For at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation For general pages on the page of	Synonyms	IVL			
SwissProt Unigene Immunogen Purified involucrin from human keratinocytes Antibody target cellular Verified antibody applications Species reactivity Antibody application notes HC (FFPE) (verified) Dog, Gorilla, Human, Monkey, Owl, Pig Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunohistology formalin-fixed 0.1-0.2 ug/mL, Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at Img/mL PBS for 5 min at 37C, enzyme digestion is better than citrate buffer based epitope unmasking, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions Tore at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Antibody research areas Antibody research areas Cancer	Human gene symbol	IVL			
Unigene 516439 Immunogen Purified involucrin from human keratinocytes Antibody target cellular Verified antibody applications Species reactivity Antibody application notes IHC (FFPE) (verified) Dog, Gorilla, Human, Monkey, Owl, Pig Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunohistology formalin-fixed 0.1-0.2 ug/mL, Staining of formalin-fixed types is enhanced by digestion with trypsin or Protease XXV at Img/mL PBS for 5 min at 37C, enzyme digestion is better than citrate buffer based epitope unmasking, Flow Cytometry 0.5-1 ug/million cells(0.1 mL, Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions Storage Conditions Tore at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Antibody research areas Antibody research areas Antibody research areas Cancer	Entrez gene ID	3713			
Immunogen	SwissProt	P07476			
Antibody target cellular Verified antibody applications Species reactivity Antibody application notes Pog. Gorilla, Human, Monkey, Owl, Pig Hicher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunohistology formalin-fixed 0.1-0.2 ug/ml., Staining of formalin-fixed 0.1-0.2 ug/ml. Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions Store at 2 to 8 ° C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 ° C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Antibody/conjugate formulation For gesearch use only (RUO) Antibody/conjugate formulation Antibody sazide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Puffled, BSA-free: 1 mg/mL in PBS upfled: 0.2 mg/mL in PBS upfle	Unigene	516439			
Verified antibody applications Species reactivity Antibody application Species reactivity Antibody application and the species of the speci	Immunogen				
Verified antibody applications Dog, Gorilla, Human, Monkey, Owl, Pig	Antibody target cellular	Cytoplasmic			
Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunohistology formalin-fixed 0.1-0.2 ug/ml., Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at 1 mg/ml. PBS for 5 min at 37C, enzyme digestion is better than citrate buffer based epitope unmasking, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Conjugates: 0.1 mg/ml. in PBS/0.1% BSA/0.05% BSA/0.05% BSA/0.05% azide, Purified: 0.2 mg/ml. in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/ml. in	Verified antibody	IHC (FFPE) (verified)			
direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunohistology formalin-fixed 0.1-0.2 ug/ml, Staining of formalin-fixed 0.1-0.2 ug/ml, Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at 1 mg/ml. PBS for 5 min at 37C, enzyme digestion is better than citrate buffer based epitope unmasking, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Immunofluorescence, Does not react with mouse; others not known, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Room temperature Storage Conditions Storage 1 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% BSA/0.05% BSA/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05%	applications Species reactivity	Dog, Gorilla, Human, Monkey, Owl, Pig			
and granular layers in normal skin. Shipping condition Room temperature Storage Conditions Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS without azide Antibody research areas Cancer	Antibody application notes	conjugates than for indirect detection with secondary antibody, which with secondary antibody, which with secondary antibody, which with secondary and the with the w			
Storage Conditions Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation SSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS Antibody research areas Cancer	Positive control				
conjugates from light, Note: store BSA-free antibodies at -10 to -35 ° C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation SAN-0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS Antibody research areas Cancer	Shipping condition	Room temperature			
date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation SA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Antibody research areas Cancer	Storage Conditions	conjugates from light, Note: store			
Antibody/conjugate formulation Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Antibody research areas Cancer	Shelf life	date of receipt when stored as			
formulation BSÅ/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Antibody research areas Cancer	Regulatory status	For research use only (RUO)			
		BSÅ/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS			
Tumor expression Squamous cell carcinoma	Antibody research areas	Cancer			
	Tumor expression	Squamous cell carcinoma			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.