Involucrin Monoclonal Mouse Antibody (IVRN/827)



Product Description

This antibody recognizes a protein of 66 kDa-170 kDa, identified as involucrin. In Western blotting of cultured human keratinocytes, this MAb reacts with a 120 kDa protein. Involucrin is expressed in a range of stratified squamous epithelia, including the cornea, which lacks a distinct cornified layer. In normal epidermis, it is first expressed in the upper spinous layers, and in keratinocyte cultures, all cells that have left the basal layer express it. Involucrin expression is altered in pathological conditions: in psoriasis and other benign epidermal hyperplasias, involucrin expression begins closer to the basal layer than normal; expression is abnormal in squamous cell carcinomas and premalignant lesions, and is reduced in severe dysplasias of the larynx and cervix. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 0827, Anti-Involucrin (IVRN/827)

Call us: 800-304-5357 Email: btinfo@biotium.com

Antibody number Antibody reactivity (target) Antibody reactivity (target) Primary Host species Mouse Clonality Monoclonal VRN/827 Isotype IgG1, kappa Molecular weight 66-170 kDa Synonyms IVL Human gene symbol VL Entrez gene ID 3713 SwissProt Unigene 516439 Immunogen Priffed involucrin from human keratinocytes Verified antibody Antibody target cellular localization Species reactivity Antibody application notes Antibody application Index on a special	Product attributes				
Host species Mouse	Antibody number	#0827			
Host species Mouse Clonality Monoclonal Clone IVRN/827 Isotype IgG1, kappa Molecular weight 66-170 kDa Synonyms IVL Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Werified antibody Antibody target cellular localization Species reactivity Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofilucrescence: 1-2 ug/mL, Immunofilustology formalin-fixed ol. 1-0.2 ug/mL, Staining of formalin-fixed ol. 1-0.2 ug/mL at 30°C, Flore the under the publication with type in the store of the publication should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Room temperature Storage Conditions Storage Conditions Storage Conditions For eact 1 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation PBS0.0.05% BSA, Duffied: 0.2 mg/mL in PBS0.05% BSA, 0.05% azide, Puffed, BSA-free: 1 mg/m		Involucrin			
Clone IVRN/827 Isotype IgG1, kappa Molecular weight 66-170 kDa Synonyms IVL Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Verified antibody Cytoplasmic Cytoplasmic Incalization Species reactivity Antibody application notes Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofluorescence: 1-2 ug/mL, Immunofluorescence: 1-2 ug/mL, Immunofluorescence: 1-2 ug/mL, Immunofluorescence: 1-2 ug/mL gestion with trypsin or Protease XXV at 1 mg/mL PBS for 5 min at 37C, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Note than enzyme digestion is better than citrate buffer based epitope unmasking, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Room temperature Storage Conditions Storage Conditions Storage Conditions For research use only (RUO) Antibody/conjugate formulation Antibody/conjugate formulation Passfree: 1 mg/mL in PBS/0.1% BSA/ 0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/ 0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/ 0.05% azide, Purified, BSA-free: 1 mg/mL in PBS urified: 0.2 mg/mL in PBS/0.05% BSA/ 0.05% azide, Purified, BSA-free: 1 mg/mL in PBS urified: 0.2 mg/mL in PBS/0.05% BSA/ 0.05% azide, Purified: 0	Antibody type	Primary			
Clone IVRN/827 Isotype IgG1, kappa Molecular weight 66-170 kDa Synonyms IVL Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Verified antibody Cytoplasmic Icoalization Species reactivity Antibody target cellular localization Species reactivity Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofluorescent 1-2 ug/mL, Immunoflu	Host species	Mouse			
Isotype IgG1, kappa	Clonality	Monoclonal			
Molecular weight Synonyms IVL	Clone	IVRN/827			
Synonyms IVL Human gene symbol IVL Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Verified antibody IHC (FFPE) (verified) Obstantial Coalization Species reactivity Antibody application notes Antibody application notes Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunofluorescence: 1-2 ug/mL, Immunofluorescence: 1-2 ug/mL, Immunofluorescence: 1-2 ug/mL, Staining of formalin-fixed of 1-0.2 ug/mL, Staining of formalin-fixed itssues is enhanced by digestion with trypsin or Protease XXV at 1 mg/mL PBS for 5 min at 37C, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Note than enzyme digestion is better than citrate buffer based epitope unmasking, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Room temperature Storage Conditions Storage Conditions Storage Conditions Storage Conditions For al 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Antibody/conjugate formulation Antibody sesearch areas Antibody research areas Antibody research areas Cancer	Isotype	IgG1, kappa			
Human gene symbol Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Verified antibody IHC (FFPE) (verified) Otyoplasmic Homan Antibody target cellular localization Species reactivity Antibody application notes Antibody application notes Otyoplasmic Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofiburoescence: 1-2 ug/mL, Immunohisology formalin-fixed 0-1-0-2 ug/mL, Staining of formalin-fixed tissues is enhanced by digestion with typsin or Protease XVV at 1 mg/mL PBS for 5 min at 37C, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Note than enzyme digestion is better than citrate buffer based epitope unmasking, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Tore at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation PBS-0.05% azide, HRP conjugates: 0.1 mg/mL in PBS-0.05% BSA, Purified: 0.2 mg/mL in PBS-0.05% BSA, Purified: 0.2 mg/mL in PBS-0.05% BSA, Purified: 0.2 mg/mL in PBS-0.05% BSA, Doustined: 0.2 mg/mL in PBS-0.05% BSA, Doustined: 0.2 mg/mL in PBS-0.05% BSA, Doustined: 0.2 mg/mL in PBS-0.05% BSA, Purified: 0.2 mg/mL in PBS-0.05% B	Molecular weight	66-170 kDa			
Entrez gene ID 3713 SwissProt P07476 Unigene 516439 Immunogen Purified involucrin from human keratinocytes Verified antibody IHC (FFPE) (verified) Antibody target cellular localization Species reactivity Antibody application notes Oytoplasmic Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunohistology formalin-fixed 0.1-0.2 ug/ml., Staining of formalin-fixed o1-0.2 ug/ml., Staining of formalin-fixed tissues is enhanced by digestion with typsin or Protease XXV at 1 mg/ml. PBS for 5 min at 37C, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Note than enzyme digestion is better than citrate buffer based epitope unmasking, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions Storage Conditions Torage Conditions Storage Conditions Storage Conditions Storage Conditions Torage Conditions Storage Conditions Storage Conditions Conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Antibody research areas Antibody research areas Antibody research areas	Synonyms	IVL			
SwissProt Unigene Immunogen Purified involucrin from human keratinocytes Verified antibody IHC (FFPE) (verified) Oytoplasmic Icoalization Species reactivity Antibody application notes Antibody application notes Purified involucrin from human keratinocytes Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofishody, Immunofishody, Immunofishody formalin-fixed 0.1-0.2 ug/ml., Staining of formalin-fixed 0.1-0.2 ug/ml., Staining of formalin-fixed to 1-0.2 ug/ml., Staining of formalin-fixed to suse is enhanced by digestion with typsin or Protease XXV at 1 mg/ml. PBS for 5 min at 37C, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Note than enzyme digestion is better than citrate buffer based epitope unmasking, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Torage Conditions Storage Conditions Storage Conditions Storage Conditions Conjugates from light, Note: stores BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation PBS/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in P	Human gene symbol	IVL			
Unigene 516439 Immunogen Purified involucrin from human keratinocytes Verified antibody Antibody target cellular localization Species reactivity Antibody application notes Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofilatody,	Entrez gene ID	3713			
Immunogen	SwissProt	P07476			
Verified antibody IHC (FFPE) (verified)	Unigene	516439			
Antibody target cellular localization Species reactivity Antibody application notes Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunoflistology formalin-fixed 0.1-0.2 ug/mL, Staining of formalin-fixed 0.1-0.2 ug/ml. pS for 5 min at 37C, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Note than enzyme digestion is better than citrate buffer based epitope unmasking, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1%, BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0	Immunogen				
Iocalization Species reactivity Human	Verified antibody	IHC (FFPE) (verified)			
Antibody application notes Antibody application notes Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofistology formalin-fixed 0.1-0.2 ug/mL, Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at 1 mg/mL PBS for 5 min at 37C, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Note than enzyme digestion is better than citrate buffer based epitope unmasking, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation The BS0.0.05% BSA, Purified: 0.2 mg/mL in PBS0.05% BSA, Purified: 0.2 mg/mL in PBS.0.05% BSA		Cytoplasmic			
direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunofilures detection with secondary and formalin-fixed 0.1-0.2 ug/mL, Staining of formalin-fixed 0.1-0.2 ug/mL, Staining of formalin-fixed 0.1-0.2 ug/mL, Staining of formalin-fixed bissues is enhanced by digestion with trypsin or Protease XXV at 1mg/mL PBS for 5 min at 37C, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Note than enzyme digestion is better than citrate buffer based epitope unmasking, Optimal dilution for a specific application should be determined by user Positive control MCF-7 cells. Localized to upper spinous and granular layers in normal skin. Shipping condition Storage Conditions Storage Conditions Storage Conditions Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% BSA/0.05% BSA/0.05% BSA/0.05% BSA/0.05% BSA/0.05% BSA/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% azide, Purified: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 2 mg/mL in PBS/0.05% BSA/0.05% azide,	Species reactivity	Human			
and granular layers in normal skin. Shipping condition Room temperature Storage Conditions Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% CSA/0.05% C		direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.1-0.2 ug/mL, Staining of formalin-fixed tissues is enhanced by digestion with trypsin or Protease XXV at 1mg/mL PBS for 5 min at 37C, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Note than enzyme digestion is better than citrate buffer based epitope unmasking, Optimal dilution for a specific application should			
Storage Conditions Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation SSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS without azide Antibody research areas Cancer	Positive control				
conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Shelf life Guaranteed for at least 24 months from date of receipt when stored as recommended Regulatory status Antibody/conjugate formulation Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.65% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Antibody research areas Cancer	Shipping condition	Room temperature			
date of receipt when stored as recommended Regulatory status For research use only (RUO) Antibody/conjugate formulation SA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Antibody research areas Cancer	Storage Conditions	conjugates from light, Note: store			
Regulatory status For research use only (RUO)	Shelf life	date of receipt when stored as			
Antibody/conjugate formulation Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide Antibody research areas Cancer	Regulatory status				
· .	Antibody/conjugate	Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS			
Tumor expression Squamous cell carcinoma	Antibody research areas	Cancer			
	Tumor expression	Squamous cell carcinoma			

Antibody # prefix BNC04	Conjugation CF®405S	Ex/Em (nm) 404/431	Laser line 405	Detection channel DAPI (microscopy), AF405	Dye Features CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.