Myeloid Associated Differentiation Marker Monoclonal Mouse Antibody (MYADM/971)



Product Description

Recognizes a myeloid associated differentiation antigen in the cytoplasm of mature granulocytes. It shows no reactivity with any other cell type in human tissues. Markers of myeloid cells are useful in the identification of different levels of cellular différentiation. It reacts with early precursor and mature forms of human and monkey myeloid cells. This MAb is useful for the detection of myeloid leukemias and granulocytic sarcomas. It can be used as a marker of granulocytes in normal tissues or inflammatory processes. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 0971, Anti-Myeloid Associated Differentiation Marker (MYADM/971)

Call us: 800-304-5357

Product attributes Antibody number Myeloid Associated Differentiation Marker Antibody reactivity (target) Antibody type Primary Host species Clonality Monoclonal Clone MYADM/971 Isotype laG1 Molecular weight Unknown MYADM; myeloid associated Synonyms differentiation marker; Myeloic upregulated protein; Protein SB135 Entrez gene ID 91663 SwissProt Q96S97 Unigene Immunogen Recombinant human MYADM protein IHC (FFPE) (verified) Verified antibody applications Antibody target cellular Cytoplasmic Species reactivity Human, Macaque Higher concentration may be required for Antibody application direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissue ugrint, Statining tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 ML, Optimal dilution for a specific application should be determined by Positive control HL60 cells. Tonsil or lymph node Shipping condition Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Storage Conditions Guaranteed for at least 24 months from date of receipt when stored as Shelf life recommended Regulatory status For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Antibody/conjugate formulation Purified, BSA-free: 1 mg/mL in PBS without azide

Antibody research areas Immunology

Granulocytes, Myeloid cells

Cell/tissue expression

Email: btinfo@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience