CD147 Monoclonal Mouse Antibody (BSG/963)



Product Description

This MAb recognizes extracellular epitope of human CD147. It is expressed more intensely on thymocytes than on mature peripheral blood T cells. CD147 is important in spermatogenesis, embryo implantation, neural network formation, and tumor progression. It stimulates the production of interstitial collagenase, gelatinase A, stromelysin-1 and various metalloproteinases (MMPs) by fibroblasts. These enzymes are important factors in cancer invasion and metastasis. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. Catalog number key for antibody number 0963, Anti-CD147 (BSG/963)

Product attributes				
Antibody number	#0963			
Antibody reactivity (target) Antibody type	CD147			
	Primary			
Host species	Mouse Monoclonal			
Clonality				
Clone	BSG/963			
Isotype	lgG1			
Molecular weight	35 kDa (non-reduced); 40 kDa (reduced)			
Synonyms	5A11 Antigen; 5F7; Basigin; Blood brain barrier HT7 antigen BSG; Extracellular Matrix Metalloproteinase Inducer (EMMPRIN); Leukocyte Activation Antigen M6; Neurothellir, OK Blood Group Antigen; TCSF; Tumor Cell-derived Collagenase Stimulatory Factor			
Human gene symbol	BSG			
Entrez gene ID	682			
SwissProt	P35613			
Unigene	501293			
Immunogen	Recombinant human BSG protein			
Antibody target cellular	Plasma membrane IHC (FFPE) (verified)			
Verified antibody				
applications Species reactivity	Human			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunchistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:100-1:200, Optimal dilution for a specific application should be determined by user			
Positive control	HSB2 cells. Renal Cell, Colon or Testicular carcinoma			
Shipping condition	Room temperature			
Storage Conditions	Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C			

Guaranteed for at least 24 months from date of receipt when stored as recommended

Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide,

For research use only (RUO)

Antibody research areas Cancer, Developmental biology

Call us: 800-304-5357 Email: btinfo@biotium.com

Product attributes

Shelf life

Regulatory status

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.