Bcl-6 Monoclonal Mouse Antibody (BCL6/850)



Product Description

Bcl-6 is a transcriptional repressor mainly required for germinal center (GC) formation and antibody affinity maturation which has different mechanisms of action specific to the lineage and biological functions. It forms complexes with different corepressors and histone deacetylases to repress the transcriptional expression of different subsets of target genes. Bcl-6 allows GC B-cells to proliferate very rapidly in response to T-cell dependent antigens and tolerate the physiological DNA breaks required for immunglobulin class switch recombination and somatic hypermutation without inducing a p53/TP53-dependent apoptotic response. Antibodies to Bcl-6 are helpful in a number of diagnostic settings, for example in the differential diagnosis of small B-cell lymphoma, and in distinguishing classical Hodgkin lymphoma from nodular lymphocyte predominant Hodgkin lymphoma (NLPHL).

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0850, Anti-Bcl-6 (BCL6/850)

Product attributes

Call us: 800-304-5357

Product attributes				
Antibody number	#0850			
Antibody reactivity (target)	Bcl-6			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	BCL6/850			
Isotype	IgG1, kappa			
Molecular weight	95 kDa			
Human gene symbol	BCL6			
Synonyms	Bcl-6; B-Cell Lymphoma 6 Protein; B-cell lymphoma 5 protein; Zinc Finger Protein 51; zinc finger transcription factor BCL6S; BCL6 Transcription Repressor			
Entrez gene ID	604			
SwissProt	P41182			
Unigene	478588			
Immunogen	Recombinant human bcl-6 protein			
Antibody target cellular localization	Nucleus			
Species reactivity	Human, Mouse			
Antibody application notes	n notes For coating for ELISA, order Ab without BSA, Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Optimal dilution and staining procedure for a specific application should be determined by user, Recommended starting concentrations for titration are 1-2 ug/mL for most applications, or 1 ug/million cells/100 uL for flow cytometry			
Shipping condition	Room temperature			
Storage Conditions	Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C			
	Trotor otoro Borr mod antibodico at Toto Go G			
Shelf life	Guaranteed for at least 24 months from date of receipt when stored as recommended			
Shelf life Regulatory status	Guaranteed for at least 24 months from date of receipt when			
	Guaranteed for at least 24 months from date of receipt when stored as recommended			
Regulatory status	Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO)			
Regulatory status Positive control Antibody/conjugate	Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Raji or Ramos cells. Tonsil or Hodgkin's lymphoma. Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in			
Regulatory status Positive control Antibody/conjugate formulation	Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Raji or Ramos cells. Tonsil or Hodgkin's lymphoma. Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

References

Garcia JF, et al. 2006. J. Histochem Cytochem. 54:31. | Pasqualucci, L., et al. 2003. Leuk. Lymphoma 44 Suppl 3: S5-12. | Ree, H.J., et al. 2003. Hum. Pathol. 34: 610-616

This datasheet was generated on December 12, 2025 at 04:58:51 AM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/monoclonal-anti-bcl-6-bcl6850/