c-Myc (EQKLISEEDL) Monoclonal Mouse Antibody (9E10.3)



Product Description

The epitope of this antibody spans the Myc tag epitope, aa 410-419 (EQKLISEEDL), which is a specific portion of an alpha helical region of human c-myc protein. Because the epitope is located at the dimerization site of the protein, the antibody is not sensitive for detecting endogenous c-myc, making it suitable for detecting Myc tagged proteins or myc overexpression. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or overexpressed in a variety of tumors. Over-expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. **Stock status**: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Call us: 800-304-5357

Product attributes				
Antibody number	#0596			
Antibody reactivity	с-Мус			
(target) Antibody type	Primary, Tag Antibody			
Host species	Mouse			
Clonality	Monoclonal			
Clone	9E10.3			
Isotype	IgG1, kappa			
Molecular weight	62-64 kDa			
Synonyms	Class E basic helix-loop-helix protein 39 (bHLHe39); MRTL; Myc2; Niard; Nird; Proto-oncogene c-Myc; RNCMYC; Transcription factor p64; Transcriptional regulator Myc-A; V-Myc avian myelocytomatosis viral oncogene homolog			
Human gene symbol	MYC			
Entrez gene ID	4609			
SwissProt	P01106			
Unigene	202453			
Immunogen	A synthetic peptide, corresponding to aa 408-439 (ASSECTION OF THE ASSECTION OF THE ASSECTI	₹NS(
Verified antibody	IHC (FFPE) (verified)			
Antibody target cellular	Nucleus			
localization Species reactivity	Human			
Antibody application notes	Immunohistology formalin-fixed 1-2 ug/ml., Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Immunofluorescence 1-2 ug/ml. Does not react with mouse, rat, or chicken, others not tested, Optimal dilution for a specific application should be determined by user			
Positive control	HL-60 cells or breast carcinoma			
Shipping condition	Room temperature			
Storage Conditions	Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C			
Shelf life	Guaranteed for at least 24 months from date of receipt when stored as recommended			
Regulatory status	For research use only (RUO)			
Antibody/conjugate formulation	Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS			

Antibody research areas Cancer, Cell cycle, Transcription factors

Email: btinfo@biotium.com

Antibody # prefix BNC04	Conjugation CF®405S	Ex/Em (nm) 404/431	Laser line 405	Detection channel DAPI (microscopy), AF405	Dye Features CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.

BSA-free