c-Myc Monoclonal Mouse Antibody (MYC909)



Product Description

This antibody recognizes a transcription factor of 64-67 kDa, identified as c-myc. This MAb shows no cross-reaction with v-myc. c-myc is involved in the control of cell proliferation and differentiation and is amplified and/or over-expressed in a variety of tumors. Over-expression of c-myc protein occurs frequently in luminal cells of prostate intraepithelial neoplasia as well as in most primary carcinomas and metastatic disease. Rearrangement of the MYC gene is found in 3% to 16% of diffuse large B-cell lymphoma (DLBCL's) and in nearly 100% of Burkitt lymphomas (BL). Identifying MYC status is important in establishing final diagnosis of DLBCL, BL, or B-cell lymphoma, with features intermediate between DLBCL and BL as well as in differential diagnoses of the lymphomas.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0909, Anti-c-Myc (MYC909)

Product attributes

Product attributes			
Antibody number	#0909		
Antibody reactivity (target)	с-Мус		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	MYC909		
Isotype	IgG1, kappa		
Molecular weight	62-64 kDa		
Synonyms	Class E basic helix-loop-helix protein 39 (bHLHe39); MRTL; Myc2; Niard; Nird; Proto-oncogene c-Myc; RNCMYC; Transcription factor 64; Transcriptional regulator Myc-A; V-Myc avian myelocytomatosis viral oncogene homolog		
Human gene symbol	MYC		
Entrez gene ID	4609		
SwissProt	P01106		
Unigene	202453		
Immunogen	Recombinant human c-myc protein		
Verified antibody applications	IHC (FFPE) (verified)		
Antibody target cellular localization	Nucleus		
Species reactivity	Human		
Antibody application notes	Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1		
	mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user		
Positive control	mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user HL-60 cells or breast carcinoma		
Positive control Shipping condition	specific application should be determined by user		
	specific application should be determined by user HL-60 cells or breast carcinoma		
Shipping condition	specific application should be determined by user HL-60 cells or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
Shipping condition Storage Conditions	specific application should be determined by user HL-60 cells or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when		
Shipping condition Storage Conditions Shelf life	specific application should be determined by user HL-60 cells or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended		
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	specific application should be determined by user HL-60 cells or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 8SA-free: 1 mg/mL in		

Chinese hamster ovary cells. Inquire for the specific lot.

Call us: 800-304-5357 Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on September 16, 2025 at 03:29:05 AM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/monoclonal-anti-c-myc-myc909/