## **Ep-CAM / CD326 Monoclonal Mouse Antibody** (VU-1D9)



## **Product Description**

This antibody reacts with the first EGF repeat in the extracellular domain of Ep-CAM. It is a 40-43 kDa transmembrane epithelial glycoprotein, also identified as epithelial specific antigen (ESA), or epithelial cellular adhesion molecule (Ep-CAM). It is expressed on baso-lateral cell surface in most simple epithelia and a vast majority of carcinomas with the exception of adult squamous epithelium, hepatocytes and gastric epithelial cells. This antibody has been used to distinguish adenocarcinoma from pleural mesothelioma and hepatocellular carcinoma. This antibody is also useful in distinguishing serous carcinomas of the ovary from mesothelioma. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

## References

Note: References for this clone sold by other suppliers may be listed for expected applications.

Mol Cell Biol 21(7): 2570-2580. (western; epitope mapping)

Product attributes				
Antibody number	#0017			
Antibody reactivity	CD326, Ep-CAM			
(target) Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	VU-1D9			
Isotype	IgG1, kappa			
Molecular weight	40-43 kDa			
Synonyms	Adenocarcinoma-associated Antigen; Cell Surface Glycoprotein Trop-1; EGP2; EGP314; EGP40; Epithelial Cell Adhesion Molecule; Epithelial Glycoprotein 314; ESA; KSA; TACD1; TROP1; Tumor-associated Calcium Signal Transducer 1 (TACSTD1)			
Human gene symbol	TACSTD1			
Entrez gene ID	4072			
SwissProt	P16422			
Unigene	542050			
Immunogen	Small cell lung carcinoma cells			
Antibody target cellular	Exosomes/EVs, Plasma membrane			
Species reactivity	Human			
Verified antibody applications	Exosome staining (verified), Flow (surface) (verified), IF (verified), IHC (FFPE) (verified)			
Expected antibody	WB (published for clone)			
applications Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunofluorescence: 1-2 ug/mL, Does not react with rat or ferret; others not known, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user			
Positive control	HT29 cells or breast tumor			
Shipping condition	Room temperature			
Storage Conditions	Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C			
Shelf life	Guaranteed for at least 24 months from date of receipt when stored as recommended			
Regulatory status	For research use only (RUO)			
Antibody/conjugate formulation	Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide			
Antibody research areas	Cancer, Cell adhesion, Exosomes/EVs			
Cell/tissue expression	Epithelial cells			
Tumor expression	Adenocarcinoma			

Call us: 800-304-5357 Email: btinfo@biotium.com

Antibody # prefix BNC04	Conjugation CF®405S	Ex/Em (nm) 404/431	<b>Laser line</b> 405	<b>Detection channel</b> DAPI (microscopy), AF405	Dye Features <a href="mailto:CF@405S Features">CF@405S Features</a>
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.