CD74 Monoclonal Mouse Antibody (CLIP/1133)



Product Description

Recognizes proteins of 33, 35 and 41 kDa, which are identified as various isoforms of CD74. Its epitope is localized in the extracellular domain of CD74. CD74 is a type II transmembrane protein which binds to the peptide binding groove of newly synthesized MHC class II alpha/beta heterodimers and prevents their premature association with endogenous polypeptides. The CD74 molecule plays a critical role in the presentation of peptides, by the MHC class II antigens, to CD4 positive lymphocytes. CD74 is expressed on MHC class II positive cells including B cells, a subset of activated T cells, monocytes, and dendritic cells and by various types of carcinomas. CD74 is expressed primarily by antigen presenting cells, such as B-lymphocytes (from before the pre-B cell stage to before the plasma cell stage), macrophages, and monocytes, and many epithelial cells. Anti-CD74 stains predominantly germinal center lymphocytes and B-cell lymphomas, but rarely T-cell lymphomas. Anti-CD74 has been shown to be useful in differentiating atypical fibroxanthoma (-) from malignant fibrous histiocytoma ().

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1133, Anti-CD74 (CLIP/1133)

Product attributes

Product attributes				
Antibody number	#1133			
Antibody reactivity (target)	CD74			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	CLIP/1133			
Isotype	IgG1, kappa			
Molecular weight	33-41 kDa			
Synonyms	CLIP; DHLAG; Gamma chain of class II antigens; HLA class II histocompatibility antigen gamma chain; HLA-DR antigene-associated invariant chain; HLADR-gamma (HLADG); la antigen-associated invariant chain; la-gamma; Major histocompatibility complex class II invariant chain; MHC HLA-DR gamma chain			
Human gene symbol	CD74			
Entrez gene ID	972			
SwissProt	P044233			
Unigene	436568			
Immunogen	Recombinant full-length human CD74 protein			
Verified antibody applications	WB (verified)			
	Plasma membrane, Perinuclear			
Antibody target cellular localization	Plasma membrane, Perinuclear			
	Plasma membrane, Perinuclear Human			
localization				
localization Species reactivity	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/ml., Flow Cytometry 0.5-1 ug/mllion cells/0.1 ml., Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should be			
localization Species reactivity Antibody application notes	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, How Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user			
localization Species reactivity Antibody application notes Positive control	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user Daudi Cells. Tonsil or lymph node.			
Species reactivity Antibody application notes Positive control Shipping condition	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user Daudi Cells. Tonsil or lymph node. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/mL, Optimal dilution for a specific application should be determined by user Daudi Cells. Tonsil or lymph node. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when			
localization Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions Shelf life	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immonstrates the secondary antibody. Immonstrates the Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1 ug/ml, Optimal dilution for a specific application should be determined by user Daudi Cells. Tonsil or lymph node. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			
localization Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/ml., Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should be determined by user Daudi Cells. Tonsil or lymph node. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% Ed. Mg/ml. in Mg/ml. in PBS/0.05% BSA/0.05% Ed. Purified: 1 mg/mL in PBS/0.05% BSA/0.05% Ed. 1 mg/mL in Mg/ml. in PBS/0.05% Ed. 1 mg/mL in 1			
Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/ml., Flow Cytometry 0.5-1 ug/million cells/0.1 ml., Western blotting 0.5-1 ug/ml., Optimal dilution for a specific application should be determined by user Daudi Cells. Tonsil or lymph node. Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide			

Call us: 800-304-5357 Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on August 5, 2025 at 02:43:39 AM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/monoclonal-anti-cd74-clip1133/