

CD59 Monoclonal Mouse Antibody (MACIF/629)



Product Description

Reacts with human CD59, a 20 kDa glycosyl phosphatidyl-inositol (GPI)-anchored cell surface protein. CD59 regulates complement-mediated cell lysis, and it is involved in lymphocyte signal transduction. This protein is a potent inhibitor of the complement membrane attack complex, whereby it binds complement C8 and/or C9 during the assembly of this complex, thereby inhibiting the incorporation of multiple copies of C9 into the complex, which is necessary for osmolytic pore formation. It inhibits formation of MAC, thus protecting cells from complement-mediated lysis. Genetic defects in GPI-anchor attachment, that cause a reduction or loss of CD59 and CD55 on erythrocytes produce the symptoms of the disease paroxysmal hemoglobinuria (PNH). This MAb is useful for study on GPI-anchored proteins, PNH and CD59 functions. CD59 is widely distributed on cells in all tissues. The expression of CD59 on erythrocytes is important for their survival.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the <u>CF® Dye Brochure</u> for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <u>order@biotium.com</u> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0629, Anti-CD59 (MACIF/629)

Call us : 800-304-5357 E

Email: techsupport@biotium.com

Product attributes

Product attributes				
Antibody number	#0629			
Antibody reactivity (target)	CD59			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	MACIF/629			
Isotype	IgG1, kappa			
Molecular weight	20 kDa			
Synonyms	20kDa homologous restriction factor (HRF20); Complement regulatory protein; Human leukocyte antigen MIC11; MAC-inhibitory protein (MACIP); Membrane attack complex inhibiton factor (MACIF); Membrane inhibitor of reactive lysis; MIRL; MSK21; Protectin; T cell activating protein			
Human gene symbol	CD59			
Entrez gene ID	966			
SwissProt	P13987			
Unigene	278573; 709466 & 710641			
Immunogen	Recombinant full-length human CD59 protein			
Verified antibody applications	IHC (FFPE) (verified)			
Antibody target cellular localization	Plasma membrane			
Species reactivity	Human			
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed tissues is enhanced by boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user			
Positive control	Daudi, CEM, K562, HPB.ALL, Jurkat, Raji, and human lymphocytes. Tonsil, Cervix or Bladder.			
Shipping condition	Room temperature			
Storage Conditions	Store at 2 to 8 $^\circ\text{C},$ Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 $^\circ\text{C}$			
Shelf life	Guaranteed for at least 24 months from date of receipt when stored as recommended			
Regulatory status	For research use only (RUO)			
Antibody/conjugate formulation	Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide			
Antibody research areas	Hematology, Immunology			
Product origin	Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot.			

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on August 24, 2025 at 07:41:57 PM. Visit product page to check for updated information before use. Product link: <u>https://biotium.com/product/monoclonal-anti-cd59-heavy-chain-of-protectin-macif629/</u>