c-Myc Monoclonal Mouse Antibody (MYC275)



Product Description

The c-Myc protein is a transcription factor, which is encoded by the c-Myc gene on human chromosome 8q24. c-Myc is commonly activated in a variety of tumor cells and plays an important role in cellular proliferation, differentiation, apoptosis and cell cycle progression. The phosphorylation of c-Myc has been investigated and previous studies have suggested a functional association between phosphorylation at Thr58/Ser62 by glycogen synthase kinase 3, cyclin dependent kinase, ERK2 and C-Jun N terminal Kinase (JNK) in cell proliferation and cell cycle regulation. Studies also have shown that c-Myc is essential for tumor cell development in vasculogenesis and angiogenesis that distribute blood throughout the cells, and which brought extensive attention in the development of new therapeutic approach for cancer treatment. Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors. **Stock status:** Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order. **Catalog** number key for antibody number 0275, Anti-c-Myc (MYC275)

Call us: 800-304-5357

Product attributes			
Antibody number	#0275		
Antibody reactivity	с-Мус		
(target) Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	MYC275		
Isotype	IgG1, kappa		
Molecular weight	62-64 kDa		
Synonyms	Class E basic helix-loop-helix protein 39 (bHL He39); MRTL; Myc2; Niard; Nird; Proto-oncogene c-Myc; RNCMYC; Transcription factor p64; Transcriptional regulator Myc-A; V-Myc avian myelocytomatosis viral oncogene homolog		
Human gene symbol	MYC		
Entrez gene ID	4609		
SwissProt	P01106		
Unigene	202453		
Immunogen	Recombinant human c-myc protein		
Verified antibody	IHC (FFPE) (verified)		
Antibody target cellular	Nucleus		
localization Species reactivity	Human		
Antibody application notes	Immunohistology formalin-fixed 1-2 ug/mL, Staining of formalin-fixed issue requires boiling tissue sections in 10 mM Tris with 1 mM EDTA, pH 9.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/millior cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user		
Positive control	HL-60 cells or breast carcinoma		
Shipping condition	Room temperature		
Storage Conditions	Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C		

Email: btinfo@biotium.com

Antibody research areas Cancer, Cell cycle, Transcription factors

Shelf life

Regulatory status

Antibody/conjugate formulation

Guaranteed for at least 24 months from date of receipt when stored as recommended

Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide,

Purified, BSA-free: 1 mg/mL in PBS

For research use only (RUO)

Antibody # prefix BNC04	Conjugation CF®405S	Ex/Em (nm) 404/431	Laser line 405	Detection channel DAPI (microscopy), AF405	Dye Features CF@405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, LI-COR Bioscience.