## Neurofilament Monoclonal Mouse Antibody (NF421 + NFL/736)



## **Product Description**

This MAb reacts with a 200 kDa and 68 kDa protein, identified as heavy and light sub-units of neurofilaments (NF-H & NF-L). Neurofilaments make up the main structural elements of axons and dendrites and are found in neurons, peripheral nerves, and sympathetic ganglion cells. Neurofilaments consist of three major subunits with molecular weights of 68 kDa (NF-L), 160 kDa (NF-M) and 200 kDa (NF-H). Anti-neurofilament stains a number of neural, neuroendocrine, and endocrine tumors. Neuromas, ganglioneuromas, gangliogliomas, ganglioneuroblastomas, and neuroblastomas stain positively for anti-neurofilament. Neurofilaments are also present in paragangliomas as well as adrenal and extra-adrenal pheochromocytomas. Carcinoids, neuroendocrine carcinomas of the skin, and oat cell carcinomas of the lung also express neurofilament.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

**Stock status:** Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0756, Anti-Neurofilament (NF421 NFL/736)

## Product attributor

Tumor expression

Product attributes			
Antibody number	#0756		
Antibody reactivity (target)	Neurofilament		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	NF421 + NFL/736		
Isotype	IgG's		
Molecular weight	200 kDa & 68 kDa		
Synonyms	NEFH; Neurofilament H; Neurofilament Heavy Polypeptide 200kDa; Neurofilament Triplet H Protein; NF-H; NF200		
Human gene symbol	NEFH		
Entrez gene ID	4744		
SwissProt	P12036		
Unigene	198760		
Immunogen	Recombinant human neurofilament heavy sub-unit (RT-97) $\&$ light sub-unit (NR-4)		
Antibody target cellular localization	Cytoskeleton		
Verified antibody applications	IHC (FFPE) (verified)		
Species reactivity	Chicken, Human, Mouse, Pig, Rat		
Antibody application notes			
	sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:25-1:50, Optimal dilution for		
Positive control	sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:25-1:50, Optimal dilution for		
Positive control Shipping condition	sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:25-1:50, Optimal dilution for a specific application should be determined by user		
	sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:25-1:50, Optimal dilution for a specific application should be determined by user  Brain, Neuroblastoma		
Shipping condition	sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:25-1:50, Optimal dilution for a specific application should be determined by user  Brain, Neuroblastoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
Shipping condition Storage Conditions	sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:25-1:50, Optimal dilution for a specific application should be determined by user  Brain, Neuroblastoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when		
Shipping condition Storage Conditions Shelf life	sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:25-1:50, Optimal dilution for a specific application should be determined by user  Brain, Neuroblastoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended		
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:25-1:50, Optimal dilution for a specific application should be determined by user  Brain, Neuroblastoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.5% BSA/0.05% BSA/0		
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1:25-1:50, Optimal dilution for a specific application should be determined by user  Brain, Neuroblastoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide		

Lung cancer, Neuroendocrine cancer

Call us: 800-304-5357 Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on August 9, 2025 at 08:07:09 PM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/monoclonal-anti-neurofilament-hl-nf421-nfl736/">https://biotium.com/product/monoclonal-anti-neurofilament-hl-nf421-nfl736/</a>