Mucin 1 / EMA / Episialin / CD227 Monoclonal Mouse Antibody (VU-2G7)



Call us: 800-304-5357

Product Description

MAb VU-2G7 reacts with MUC1, a large transmembrane glycoprotein expressed on the ductal surface of normal glandular epithelia. The dominant epitope of VU-2G7 has not been established with epitope fingerprinting. The extracellular domain of MUC1 largely consists of a highly conserved, O-glycosylated 20 amino acids tandem repeat which can occur 30-100 times per molecule depending on the length of the allele involved. In the vast majority of human carcinomas this protein is up regulated and poorly glycosylated and appears on the cell surface in a non-polarized fashion.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0630, Anti-CD227 (VU-2G7)

References

Note: References for this clone sold by other suppliers may be listed for expected applications.

- 1. ACS Omega (2017) 2 (11), 7493-7505. (ELISA, Flow)
- 2. Arch Dermatol Res (2003) 295:146-154. (immunohistochemistry)

Product attributes				
Antibody number	#0630			
Antibody reactivity (target)	CD227, EMA, Episialin, Mucin 1			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clonality	VU-2G7			
Isotype	IgG1, kappa			
Molecular weight	265-400 kDa			
Synonyms	Breast carcinoma-associated antigen DF3; CA15-3; Carcinoma-associated mucin Episialin; Epithelial Membrane Antigen; H23AG; KL-6; MAM6; MUC-1; MUC-1/SEC; MUC-1/X; MUC1-alpha; MUC1-beta; MUC1-CT; MUC1-NT; MUC1/ZD; Mucin 1 cell surface associated; Mucin-1 subunit beta; Peanut-reactive urinary mucin; PEM; PEMT; Polymorphic epithelial mucin; PUM; Tumor-associated epithelial membrane antigen			
Human gene symbol	MUC1			
Entrez gene ID	4582			
SwissProt	P15941			
Unigene	89603			
Immunogen	Synthetic glycosylated MUC1 60mer tandem repeat NH2-(HGVTSAPDT(GalNAc)RPAPGSTACOOH, conjugated to bovine serum albumin			
Antibody target cellular	Plasma membrane			
Expected antibody applications	ELISA (published for clone), Flow, surface (published for clone), IHC (FFPE) (published for clone)			
Species reactivity	Human			
Antibody application notes	Immunohistology formalin-paraffin 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Optimal dilution for a specific application should be determined by user			
Positive control	MCF-7 or MDA-231 cells. Breast, colon, ovarian, endometrial carcinoma.			
Shipping condition	Room temperature			
Storage Conditions	Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C			
Shelf life	Guaranteed for at least 24 months from date of receipt when stored as recommended			
Regulatory status	For research use only (RUO)			
Antibody/conjugate formulation	Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide,			

Purified, BSA-free: 1 mg/mL in PBS without azide

Epithelial cells, Mammary gland

Cancer, Mucins

Breast cancer

Antibody research areas

Cell/tissue expression

Tumor expression

Email: btinfo@biotium.com

Antibody # prefix BNC04	Conjugation CF®405S	Ex/Em (nm) 404/431	Laser line 405	Detection channel DAPI (microscopy), AF405	Dye Features CF®405S Features
BNC88	CF®488A	490/515	488	GFP. FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, of LI-COR Bioscience.

