NGF Receptor Monoclonal Mouse Antibody (NTR/912)



Product Description

This antibody recognizes a glycoprotein of 75 kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0912, Anti-NGFR (NTR/912)

Product attributes

Call us: 800-304-5357

Antibody number	#0912		
Antibody reactivity (target)	NGFR		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	NTR/912		
Isotype	IgG1, kappa		
Molecular weight	75 kDa		
Synonyms	CD271; Gp80-LNGFR; Low affinity nerve growth factor recepto Low affinity neurotrophin receptor p75NTR; Nerve growth factor receptor (NGFR); p75 ICD; p75 Neurotrophin receptor; Tumor necrosis factor receptor superfamily member 16 (TNFRSF16)		
Human gene symbol	NGFR		
Entrez gene ID	4804		
SwissProt	P08138		
Unigene	415768 & 681726		
Immunogen	Recombinant human p75 NGFR protein		
Antibody target cellular localization	Plasma membrane		
Species reactivity	Human, Non-human primates		
Verified antibody applications	IHC (FFPE) (verified)		
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody. Immunofluorescence: 1-2 ug/mL. Does not		

Email: techsupport@biotium.com

Shipping condition	Room temperature			
Storage Conditions	Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C			
Shelf life	Guaranteed for at least 24 months from date of receipt when stored as recommended			
Regulatory status	For research use only (RUO)			
Antibody research areas	Cancer, Developmental biology, Signal transduction			
Cell/tissue expression	Neural crest cells			
Tumor expression	Melanoma, Neuroendocrine cancer			
Antibody/conjugate formulation	Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/m in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/ml PBS without azide			
Positive control	Melanoma, Neuronal axons, Schwann cells, and perineural cells of peripheral nerves, Soma and axons of sensory neurons, and ganglionic satellite cells, Tumors of nerve sheath differentiation,			

react with mouse or rat, others not known, Immunohistology

requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a

formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissu

specific application should be determined by use

Schwannoma, Neurofibroma

Product origin

Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot.

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on July 12, 2025 at 02:46:03 AM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/monoclonal-anti-ngfr-nerve-growth-factor-receptor-ntr912/