## NGF Receptor Monoclonal Mouse Antibody (NTR/912)



## **Product Description**

This antibody recognizes a glycoprotein of 75 kDa, identified as low affinity Nerve Growth Factor (NGF) Receptor (p75NGFR) or Neurotrophin Receptor (p75NTR). NGFR is expressed in various neural crest cells and their tumors such as melanocytes, melanomas, neuroblastomas, pheochromocytomas and neurofibromas. Reportedly, anti-NGFR is a reliable marker for desmoplastic and neurotropic melanomas. NGFR is expressed in mature non-neural cells such as perivascular cells, dental pulp cells, lymphoidal follicular dendritic cells, basal epithelium of oral mucosa and hair follicles, prostate basal cells, and myoepithelial cells. Anti-NGFR stains the myoepithelial cells of breast ducts and intra-lobular fibroblasts of breast ducts.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0912, Anti-NGFR (NTR/912)

## Call us: 800-304-5357

Shipping condition

| Product attributes                    |  |  |  |
|---------------------------------------|--|--|--|
| Antibody number                       | #0912  |  |  |
| Antibody reactivity (target)          | NGFR   |  |  |
| Antibody type                         | Primary  |  |  |
| Host species                          | Mouse  |  |  |
| Clonality                             | Monoclonal   |  |  |
| Clone                                 | NTR/912  |  |  |
| Isotype                               | IgG1, kappa  |  |  |
| Molecular weight                      | 75 kDa   |  |  |
| Synonyms                              | CD271; Gp80-LNGFR; Low affinity nerve growth factor receptor;<br>Low affinity neurotrophin receptor p75NTR; Nerve growth factor<br>receptor (NGFR); p75 ICD; p75 Neurotrophin receptor; Tumor<br>necrosis factor receptor superfamily member 16 (TNFRSF16) |  |  |
| Human gene symbol                     | NGFR   |  |  |
| Entrez gene ID                        | 4804   |  |  |
| SwissProt                             | P08138   |  |  |
| Unigene                               | 415768 & 681726  |  |  |
| Immunogen                             | Recombinant human p75 NGFR protein   |  |  |
| Antibody target cellular localization | Plasma membrane  |  |  |
| Species reactivity                    | Human, Non-human primates  |  |  |
| Verified antibody applications        | IHC (FFPE) (verified)  |  |  |
| Antibody application notes            | Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 1-2 ug/mL, Does not   |  |  |

| secondary antibody, Immunofluorescence: 1-2 ug/mL, Does not react with mouse or rat, others not known, Immunohistology formalin-fixed 0.5-1 ug/mL, Staining of formalin-fixed tissues requires boiling tissue sections in 10 mM citrate buffer, pH 6.0, for 10-20 min followed by cooling at RT for 20 minutes, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user |
|--|
| Room temperature   |
| Store at 2 to 8 °C. Protect fluorescent conjugates from light.   |

| Storage Conditions             | Store at 2 to 8 $^{\circ}$ C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 $^{\circ}$ C   |  |  |
|--------------------------------|---|--|--|
| Shelf life                     | Guaranteed for at least 24 months from date of receipt when stored as recommended   |  |  |
| Regulatory status              | For research use only (RUO)   |  |  |
| Antibody research areas        | Cancer, Developmental biology, Signal transduction  |  |  |
| Cell/tissue expression         | Neural crest cells  |  |  |
| Tumor expression               | Melanoma, Neuroendocrine cancer   |  |  |
| Antibody/conjugate formulation | Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide                       |  |  |
| Positive control               | Melanoma, Neuronal axons, Schwann cells, and perineural cells of peripheral nerves, Soma and axons of sensory neurons, and ganglionic satellite cells, Tumors of nerve sheath differentiation, Schwannoma, Neurofibroma |  |  |
| Product origin                 | Product may contain either bovine serum albumin (BSA) from  |  |  |

origin

Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in Chinese hamster ovary cells. Inquire for the specific lot.

| Antibody # prefix | Conjugation           | Ex/Em (nm) | Laser line | Detection channel           | Dye Features     |
|-------------------|-----------------------|------------|------------|-----------------------------|------------------|
| BNC04             | CF®405S               | 404/431    | 405        | DAPI (microscopy),<br>AF405 | CF®405S Features |
| BNC88             | CF®488A               | 490/515    | 488        | GFP, FITC                   | CF®488A Features |
| BNC68             | CF®568                | 562/583    | 532, 561   | RFP, TRITC                  | CF®568 Features  |
| BNC94             | CF®594                | 593/614    | 561        | Texas Red®                  | CF®594 Features  |
| BNC40             | CF®640R               | 642/662    | 633-640    | Cy®5                        | CF®640R Features |
| BNC47             | CF®647                | 650/665    | 633-640    | Cy®5                        | CF®647 Features  |
| BNC74             | CF®740                | 742/767    | 633-685    | 775/50                      | CF®740 Features  |
| BNCB              | Biotin                | N/A        | N/A        | N/A                         |                  |
| BNUB              | Purified              | N/A        | N/A        | N/A                         |                  |
| BNUM              | Purified,<br>BSA-free | N/A        | N/A        | N/A                         |                  |

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on December 12, 2025 at 05:40:53 AM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/monoclonal-anti-ngfr-nerve-growth-factor-receptor-ntr912/">https://biotium.com/product/monoclonal-anti-ngfr-nerve-growth-factor-receptor-ntr912/</a>