## S100A9 Monoclonal Mouse Antibody (47-8D3)

## **Product Description**

This MAb stains the cytoplasm of macrophages and histiocytes in hematopoietic organs, Kupffer's cells of the liver and Langerhan's cells of the skin. It also stains the mantle zone B-lymphocytes of the lymph node and spleen, spermatogonia, and chief cells of the stomach. S100A9 is expressed by macrophages in acutely inflamed tissues and in chronic inflammation. It is detected in peripheral blood leukocytes, in neutrophils and granulocytes. It is present at sites of vascular inflammation. S100A9 is also expressed in epithelial cells constitutively or induced during dermatoses. S100A9 is a Calcium-binding protein. It has antimicrobial activity towards bacteria and fungi. It is important for resistance to invasion by pathogenic bacteria. It up-regulates transcription of genes that are under the control of NF-kappa-B. S100A9 plays a role in the development of endotoxic shock in response to bacterial lipopolysaccharide (LPS). It promotes tubulin polymerization when unphosphorylated. It also promotes phagocyte migration and infiltration of granulocytes at sites of wounding. It plays a role as a pro-inflammatory mediator in acute and chronic inflammation and up-regulates the release of IL8 and cell-surface expression of ICAM1.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1093, Anti-S100A9 (47-8D3)

## Product attributes

Product attributes				
Antibody number	#1093			
Antibody reactivity (target)	S100A9			
Antibody type	Primary			
Host species	Mouse			
Clonality	Monoclonal			
Clone	47-8D3			
Isotype	IgG1, kappa			
Molecular weight	14 kDa			
Synonyms	S100-A9; MRP14; calgranulin-B; calprotectin L1H subunit; leukocyte L1 complex heavy chain; migration inhibitory factor-related protein 14; S100 calcium-binding protein A9 (calgranulin B)			
Human gene symbol	S100A9			
Entrez gene ID	6280			
SwissProt	P06702			
Unigene	112405			
Immunogen	Human peripheral blood monocyte components			
Antibody target cellular localization	Cytoplasmic, Nucleus			
Species reactivity	Human, Rat			
	WB (published for clone)			
Expected antibody applications	WB (published for clone)			
	WB (published for clone)  Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofiluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at Tmg/mL PBS, 15 min at RT, Flow Cytometry 0.5-1 ug/million cells/0.1 mL			
applications	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at 1mg/mL PBS, 15 min at RT, Flow Cytometry 0.5-1 ug/million			
applications Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofiluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at 1mg/mL PBS, 15 min at RT, Flow Cytometry 0.5-1 ug/million cells/0.1 mL			
applications Antibody application notes Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofiluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at 1mg/mL PBS, 15 min at RT, Flow Cytometry 0.5-1 ug/million cells/0.1 mL.  Liver or histiocytoma			
applications Antibody application notes  Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at 1mg/mL PBS, 15 min at RT, Flow Cytometry 0.5-1 ug/million cells/0.1 mL  Liver or histiocytoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
applications Antibody application notes  Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at 1mg/mL PBS, 15 min at RT, Flow Cytometry 0.5-1 ug/million cells/0.1 mL.  Liver or histiocytoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when			
applications Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at 1 mg/mL PBS, 15 min at RT, Flow Cytometry 0.5-1 ug/million cells/0.1 mL.  Liver or histicocytoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended			
applications Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at 1mg/mL PBS, 15 min at RT, Flow Cytometry 0.5-1 ug/million cells/0.1 mL  Liver or histiocytoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; BSA-free: 1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; DSA-free: 1 mg/mL in PBS/0.05			
applications Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.5-1 ug/mL, For staining of formalin-fixed, paraffin-embedded tissues, digest with trypsin at 1mg/mL PBS, 15 min at RT, Flow Cytometry 0.5-1 ug/million cells/0.1 mL.  Liver or histiocytoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide			

Call us: 800-304-5357 Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

## References

Note: References for this clone sold by other suppliers may be listed for expected applications.

J Invest Dermatol (2010) 130(10) 2423-2430. (WB)

This datasheet was generated on September 11, 2025 at 02:37:37 PM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/monoclonal-calprotectinmrp14s100a9calgranulin-b-47-8d3/">https://biotium.com/product/monoclonal-calprotectinmrp14s100a9calgranulin-b-47-8d3/</a>