CD3e Polyclonal Rabbit Antibody (PAb)

Product Description

The immunogen for this Mab is derived from the cytoplasmic domain of CD3 epsilon chain. CD3 consists of five different polypeptide chains (designated as gamma, delta, epsilon, zeta, and eta) with MW ranging from 16-28 kDa. The CD3 complex is closely associated at the lymphocyte cell surface with the T cell antigen receptor (TCR). Reportedly, CD3 complex is involved in signal transduction to the T cell interior following antigen recognition. The CD3 antigen is first detectable in early thymocytes and probably represents one of the earliest signs of commitment to the T cell lineage. In cortical thymocytes, CD3 is predominantly intra-cytoplasmic. However, in medullary thymocytes, it appears on the T cell surface. CD3 antigen is a highly specific marker for T cells, and is present in majority of T cell neoplasms.

Also view our <u>CD3e Monoclonal Mouse Biotium Choice Antibody (SK7)</u>. The antibody belongs to the Biotium Choice list of select antibodies that have been developed and validated in-house for optimal performance. The antibody has been validated in flow cytometry.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0385, Anti-CD3e (PAb)

Product attributes

Call us: 800-304-5357

Product attributes				
Antibody number	#0385			
Antibody reactivity (target)	CD3e			
Antibody type	Primary			
Host species	Rabbit			
Clonality	Polyclonal			
Clone	PAb			
Isotype	IgG			
Molecular weight	20 kDa			
Synonyms	CD3 epsilon; CD3 TCR complex; T cell antigen receptor complex epsilon subunit of T3; T-cell surface antigen T3/Leu-4 epsilon chain; T-cell surface glycoprotein CD3 epsilon chain; T3E; TCRE; TiT3 complex			
Human gene symbol	CD3E			
Entrez gene ID	916			
SwissProt	P07766			
Unigene	3003			
Immunogen	Synthetic peptide corresponding to amino acids 156-168 (ERPPPVPNPDYEP) of the cytoplasmic domain of human CD3e			
Verified antibody applications	WB (verified)			
	Plasma membrane			
Antibody target cellular localization	Plasma membrane			
	Plasma membrane Human			
localization				
localization Species reactivity	Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Predicted to show broad species reactivity, Optimal dilution for a specific			
localization Species reactivity Antibody application notes	Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Predicted to show broad species reactivity, Optimal dilution for a specific application should be determined by user			
localization Species reactivity Antibody application notes Positive control	Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Predicted to show broad species reactivity, Optimal dilution for a specific application should be determined by user Jurkat cells. Tonsil and lymph node			
localization Species reactivity Antibody application notes Positive control Shipping condition	Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Predicted to show broad species reactivity, Optimal dilution for a specific application should be determined by user Jurkat cells. Tonsil and lymph node Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,			
localization Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions	Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Predicted to show broad species reactivity, Optimal dilution for a specific application should be determined by user Jurkat cells. Tonsil and lymph node Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when			
localization Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions Shelf life	Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Predicted to show broad species reactivity, Optimal dilution for a specific application should be determined by user Jurkat cells. Tonsil and lymph node Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended			
localization Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, immunofluorescence 1-2 ug/mL, Predicted to show broad species reactivity, Optimal dilution for a specific application should be determined by user Jurkat cells. Tonsil and lymph node Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA, Purified; 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; 10.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; 10.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; 10.2 mg/mL in			
localization Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, immunofluorescence 1-2 ug/mL, Predicted to show broad species reactivity, Optimal dilution for a specific application should be determined by user Jurkat cells. Tonsil and lymph node Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified. BSA-free: 1 mg/mL in PBS without azide			
localization Species reactivity Antibody application notes Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Antibody research areas	Human Western blotting 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Immunofluorescence 1-2 ug/mL, Predicted to show broad species reactivity, Optimal dilution for a specific application should be determined by user Jurkat cells. Tonsil and lymph node Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA/0.05% azide, Purified BSA-free: 1 mg/mL in PBS without azide Immunology Product may contain either bovine serum albumin (BSA) from bovine serum (Bos taurus), or recombinant BSA produced in			

Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on August 13, 2025 at 01:27:10 AM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/monoclonal-anti-cd3e-rabbit-pab/