

## Cytokeratin 8/18 Monoclonal Mouse Antibody (B22.1 + B23.1)



## **Product Description**

Cytokeratin 8 (CK8) belongs to the type II (or B or basic) subfamily of high molecular weight cytokeratins and exists in combination with cytokeratin 18 (CK18). This MAb cocktail recognizes all simple epithelia including glandular epithelium, for example thyroid, female breast, gastrointestinal tract, respiratory tract, and urogenital tract including transitional epithelium. All adenocarcinomas and most squamous carcinomas are positive but keratinizing squamous carcinomas are usually negative. This antibody is useful in demonstrating the presence of Paget cells; there is very little keratin 18 in the normal epidermis so only Paget cells are stained. Immunohistochemical staining with this MAb is indistinguishable from that obtained with monoclonal antibody 5D3.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 1221, Anti-Cytokeratin 8 (B22.1 B23.1)

## Call us : 800-304-5357 Email: techsupport@biotium.com

Product attributes					
Antibody number	#1221				
Antibody reactivity (target)	Cytokeratin 8/18				
Antibody type	Primary				
Host species	Mouse				
Clonality	Monoclonal				
Clone	B22.1 + B23.1				
Isotype	lgG's				
Molecular weight	52.5 kDa (CK8); 45 kDa (CK18)				
Synonyms	CARD2; CK8; CYK8; CYKER; Cytokeratin Endo A; DreK8; EndoA; K2C8; K8; Keratin 8; Krt 2.8; KRT8; Type-II Keratin Kb8				
Human gene symbol	KRT8 & KRT18				
Entrez gene ID	3856 (CK8); 3875 (CK18)				
SwissProt	P05787 (CK8); P05783 (CK18)				
Unigene	533782 & 708445 (CK8); 406013 (CK18)				
Immunogen	Cytoskeletal preparation from HeLa epitheliod carcinoma cells (B22.1) and PMC-42 breast carcinoma cells of human origin (B23.1)				
Antibody target cellular localization	Cytoskeleton				
Verified antibody applications	IHC (FFPE) (verified)				
Species reactivity	Human				
Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with				
	secondary antibody, Immunofluorescence: 1-2 ug/mL, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0 ug/mL				
Positive control	Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0				
Positive control Shipping condition	Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0 ug/mL MCF-7, HeLa, PMC-42 or A431 cells. Skin, Colon, lung or breast				
	Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0 ug/mL MCF-7, HeLa, PMC-42 or A431 cells. Skin, Colon, lung or breast carcinoma				
Shipping condition	Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0 ug/mL MCF-7, HeLa, PMC-42 or A431 cells. Skin, Colon, lung or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light,				
Shipping condition Storage Conditions	Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0 ug/mL MCF-7, HeLa, PMC-42 or A431 cells. Skin, Colon, lung or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when				
Shipping condition Storage Conditions Shelf life	Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0 ug/mL MCF-7, HeLa, PMC-42 or A431 cells. Skin, Colon, lung or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended				
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0 ug/mL MCF-7, HeLa, PMC-42 or A431 cells. Skin, Colon, lung or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS/0.05% azide, Purified, SA-free: 1 mg/mL in PBS/0.05% azide, Pu				
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0 ug/mL MCF-7, HeLa, PMC-42 or A431 cells. Skin, Colon, lung or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified, BSA-free: 1 mg/mL in PBS without azide				
Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation Antibody research areas	Cytometry 0.5-1 ug/million cells/0.1 mL, Western blotting 0.5-1.0 ug/mL MCF-7, HeLa, PMC-42 or A431 cells. Skin, Colon, lung or breast carcinoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purified, BSA-free: 1 mg/mL in PBS/0.05% BSA, Purified; D.2 mg/mL in PBS/0.05% BSA, Purified; D.2 mg/mL in PBS/0.05% azide, PURIFIEd;				

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on September 11, 2025 at 01:14:06 PM. Visit product page to check for updated information before use. Product link: https://biotium.com/product/monoclonal-anti-cytokeratin-818-b22-1-b23-1/