## PLAP Monoclonal Mouse Antibody (ALP/870)



## **Product Description**

Reacts with a 70 kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

**Stock status:** Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email <a href="mailto:order@biotium.com">order@biotium.com</a> to inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0870, Anti-PLAP (ALPP/870)

## **Product attributes**

Call us: 800-304-5357

Froduct attributes			
Antibody number	#0870		
Antibody reactivity (target)	PLAP		
Antibody type	Primary		
Host species	Mouse		
Clonality	Monoclonal		
Clone	ALP/870		
Isotype	IgG2b, kappa		
Molecular weight	70 kDa		
Synonyms	Alkaline phosphatase placental type, Alkaline phosphatase Regan Isozyme, ALP, Alp1, ALPP, Germ-cell alkaline phosphatase, nagao Isozyme, PALP, Placental alkaline phosphatase 1, placental heat-stable alkaline phosphatase, PLAP-1, PLAP1		
Human gene symbol	ALPP		
Entrez gene ID	250		
SwissProt	P05187		
Unigene	284255		
Immunogen	Recombinant full-length human ALPP protein		
Antibody target cellular localization	Plasma membrane		
Verified antibody applications	IHC (FFPE) (verified), WB (verified)		
• •			
Species reactivity	Human		
• •	Human  Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0 5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
Species reactivity	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be		
Species reactivity Antibody application notes	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
Species reactivity Antibody application notes  Positive control	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  HepG2 cells. Placenta or seminoma		
Species reactivity Antibody application notes  Positive control Shipping condition	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  HepG2 cells. Placenta or seminoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light,		
Species reactivity Antibody application notes  Positive control Shipping condition Storage Conditions	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/ml., Immunoflistology formalin-fixed 0.25-0.5 ug/ml., No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  HepG2 cells. Placenta or seminoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when		
Species reactivity Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  HepG2 cells. Placenta or seminoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended		
Species reactivity Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  HepG2 cells. Placenta or seminoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA/0.05% azide, Purified: 0.2 mg/mL in PBS/0.05% azide, Purifi		
Species reactivity Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  HepG2 cells. Placenta or seminoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% BSA-free: 1 mg/mL in PBS without azide		
Species reactivity Antibody application notes  Positive control Shipping condition Storage Conditions Shelf life Regulatory status Antibody/conjugate formulation  Cell/tissue expression	Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user  HepG2 cells. Placenta or seminoma  Room temperature  Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C  Guaranteed for at least 24 months from date of receipt when stored as recommended  For research use only (RUO)  Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.05% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA/0.05% azide, Purified; DSS without azide  Placenta		

Email: techsupport@biotium.com

Antibody # prefix	Conjugation	Ex/Em (nm)	Laser line	Detection channel	Dye Features
BNC04	CF®405S	404/431	405	DAPI (microscopy), AF405	CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNC74	CF®740	742/767	633-685	775/50	CF®740 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified, BSA-free	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, and Odyssey are registered trademarks of LI-COR Bioscience.

This datasheet was generated on September 11, 2025 at 11:25:48 AM. Visit product page to check for updated information before use. Product link: <a href="https://biotium.com/product/monoclonal-anti-plap-placental-alkaline-phosphatase-alp870/">https://biotium.com/product/monoclonal-anti-plap-placental-alkaline-phosphatase-alp870/</a>