PLAP Monoclonal Mouse Antibody (ALP/870)



Product Description

Reacts with a 70 kDa membrane-bound isozyme (Regan and Nagao type) of Placental Alkaline Phosphatase (PLAP) occurring in the placenta during the 3rd trimester of gestation. It is highly specific for PLAP and shows no cross-reaction with other isozymes of alkaline phosphatase. Anti-PLAP reacts with germ cell tumors and can discriminate between these and other neoplasms. Somatic neoplasms e.g. breast, gastrointestinal, prostatic, and urinary cancers may also immunoreact with antibodies to PLAP. Anti-PLAP positivity in conjunction with anti-keratin negativity favors seminoma over carcinoma. Germ cell tumors are usually anti-keratin positive, but they regularly fail to stain with anti-EMA, whereas most carcinomas stain with anti-EMA. Anti-PLAP has been useful in the diagnosis of gestational trophoblastic disease.

Primary antibodies are available purified, or with a selection of fluorescent CF® dyes and other labels. CF® dyes offer exceptional brightness and photostability. See the CF® Dye Brochure for more information. Note: Conjugates of blue fluorescent dyes like CF®405S and CF®405M are not recommended for detecting low abundance targets, because blue dyes have lower fluorescence and can give higher non-specific background than other dye colors.

Stock status: Because Biotium offers a large number of antibody and conjugation options, primary antibody conjugates may be made to order. Typical lead times are up to one week for CF® dye and biotin conjugates, and up to 2-3 weeks for fluorescent protein and enzyme conjugates. Please email order@biotium.com to

inquire about stock status and lead times before placing your order.

Catalog number key for antibody number 0870, Anti-PLAP (ALPP/870)

Tumor expression

Call us: 800-304-5357

#0870		
PLAP		
Primary		
Mouse		
Monoclonal		
ALP/870		
IgG2b, kappa		
70 kDa		
Alkaline phosphatase placental type, Alkaline phosphatase Regan Isozyme, ALP, Alp1, ALPP, Germ-Cell alkaline phosphatase, nagao Isozyme, PALP, Placental alkaline phosphatase 1, placental heat-stable alkaline phosphatase, PLAP-1, PLAP1		
ALPP		
250		
P05187		
284255		
Recombinant full-length human ALPP protein		
Plasma membrane		
IHC (FFPE) (verified), WB (verified)		
IHC (FFPE) (verified), WB (verified) Human		
Human		
Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofiluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application		
Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user		
Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user HepG2 cells. Placenta or seminoma		
Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user HepG2 cells. Placenta or seminoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store		
Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/mllion cells/0.1 mL, Optimal dilution for a specific application should be determined by user HepG2 cells. Placenta or seminoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as		
Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user HepG2 cells. Placenta or seminoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Guaranteed for at least 24 months from date of receipt when stored as recommended		
Human Higher concentration may be required for direct detection using primary antibody conjugates than for indirect detection with secondary antibody, Immunofluorescence: 0.5-1 ug/mL, Immunohistology formalin-fixed 0.25-0.5 ug/mL, No special pretreatment is required for the immunohistochemical staining of formalin-fixed tissues, Flow Cytometry 0.5-1 ug/million cells/0.1 mL, Optimal dilution for a specific application should be determined by user HepG2 cells. Placenta or seminoma Room temperature Store at 2 to 8 °C, Protect fluorescent conjugates from light, Note: store BSA-free antibodies at -10 to -35 °C Gauranteed for at least 24 months from date of receipt when stored as recommended For research use only (RUO) Conjugates: 0.1 mg/mL in PBS/0.1% BSA/0.65% azide, HRP conjugates: 0.1 mg/mL in PBS/0.05% BSA, Purified: 0.2 mg/mL in PBS/0.05% BSA, Purif		

Ovarian cancer, Testicular cancer

Email: btinfo@biotium.com

Antibody # prefix BNC04	Conjugation CF®405S	Ex/Em (nm) 404/431	Laser line 405	Detection channel DAPI (microscopy), AF405	Dye Features CF®405S Features
BNC88	CF®488A	490/515	488	GFP, FITC	CF®488A Features
BNC68	CF®568	562/583	532, 561	RFP, TRITC	CF®568 Features
BNC94	CF®594	593/614	561	Texas Red®	CF®594 Features
BNC40	CF®640R	642/662	633-640	Cy®5	CF®640R Features
BNC47	CF®647	650/665	633-640	Cy®5	CF®647 Features
BNCB	Biotin	N/A	N/A	N/A	
BNUB	Purified	N/A	N/A	N/A	
BNUM	Purified,	N/A	N/A	N/A	

Alexa Fluor, Pacific Blue, Pacific Orange, and Texas Red are trademarks or registered trademarks of Thermo Fisher Scientific; Cy is a registered trademark of Cytiva; IRDye, LI-COR, of LI-COR Bioscience.